


1900—1911 The Pre-Discovery Period

1900 J Elster and H Geitel, CTR Wilson: Explanation of why air becomes conductive



Elster and Geitel in their private lab

Following the discovery of radioactivity and X-rays, the ionisation of gases was systematically investigated. For these measurements, electrometers originally invented in 1789 to study electrical phenomena were used.

Julius Elster and Hans Geitel (Wolfenbüttel), and Charles Thomson Rösser Wilson (Cambridge) found that ion production appears in closed and isolated detectors even in the absence of a source. The general conclusion was that this unexpected ionisation is caused by radioactive substances in the detector's walls or in its neighbourhood. Wilson was the first to ask the question: can this penetrating radiation be extraterrestrial?

1901 CTR Wilson: Extraterrestrial radiation mentioned for the first time

1902—1903 F Linke: First investigation of penetrating radiation with balloon flights

During six balloon flights, Franz Linke (Berlin), a meteorologist and geologist, performed ionisation measurements with an electrometer provided by Elster and Geitel. He measured the ionisation compared to on the ground. It was about the same between altitudes of 1000m to 3000m and larger by a factor of four at 5500m. His published results have obviously (and unfortunately) never been recognised.



The Phoenix balloon (Berlin) in 1894, drawn by H Groß


1902—1909 General consensus that ionisation is caused by the natural radioactivity of the Earth

Over the following years, the ionisation effect was studied by many scientists in Canada, Germany, Britain and the US. The electrometers were surrounded by such different materials as water, wood, bricks and lead. The goal was to study the absorption of penetrating beta rays and gamma rays coming from radioactive elements in the ground and in the air of the environment. With increasing absorber thickness, the ionisation could be reduced, but a small fraction remained.

1908 T Wulf: Improvement of the electrometer's design



Theodor Wulf



Two-string electrometer designed by Wulf

Theodor Wulf, a German Jesuit, studied physics in Innsbruck and Göttingen. He improved the electrometer's design by replacing the thin metal foils with two thin metal fibres. This new electrometer was much easier to calibrate and became the state-of-the-art detector for outdoor measurements.

1908 A Gockel and T Wulf coin the term "cosmic radiation"

Wulf and Albert Gockel (Switzerland) were the first to study the ionisation rate on high mountains in the Alps. They did not observe strong deviations from measurements at sea level. In their publication, the term "cosmic radiation" was used for the first time.

1909 K Bergwitz, A Gockel: New high-altitude measurements with balloons



Albert Gockel

Near Braunschweig, Karl Bergwitz reached an altitude of 1300m with a balloon. He measured the ionisation with an improved electrometer as designed by Wulf. The measurements were continued by Albert Gockel (Fribourg). Neither observed a decrease of ionisation with distance from the ground, as had been expected for radiation sources in the Earth's crust. Because of problems with the detectors, it was difficult to draw definitive conclusions.



Karl Bergwitz

1910 T Wulf: Ionisation measurement on the Eiffel Tower

Wulf began a series of measurements on top of the Eiffel Tower. Even at only 300m above ground, he observed a smaller reduction in radiation when compared with theoretical estimates. He came to the same conclusion as Bergwitz and Gockel: the radioactivity of the air must contribute essentially to the measured ionisation.

1910—1911 D Pacini, GC Simpson and CS Wright: Measurement of ionising radiation on and below a water surface



Domenico Pacini with an electrometer

In Italy, Domenico Pacini (Bari) performed ionisation measurements in the bay of Livorno at a distance of 300m from the coast on the water's surface and 3m below it. Since ionising radiation from solid materials could be excluded on the sea, he concluded that the source of ionisation must be a penetrating radiation in the atmosphere.


George Simpson and Charles Wright were scientific members of RF Scott's British Antarctic Expedition. Sailing from Britain to New Zealand, they measured the ionisation and observed a dependence of the rate on the barometric air pressure, not knowing that it was connected with cosmic rays.

1912 Discovery by Victor Hess

1911

CTR Wilson:
Development of the cloud chamber and publication of the first pictures

In 1895 CTR Wilson started investigating cloud formation in dust-free air. He discovered that condensed bubbles appear when air molecules are ionised by X-rays. In 1911 Wilson demonstrated with a cloud chamber that alpha and beta rays could be visualised. Two of the published pictures contained straight tracks which were probably the first photographs of cosmic particles. One year before their discovery, Wilson misinterpreted these tracks as beta rays.



Original Wilson cloud chamber (Cavendish Museum)

1911–1912

VF Hess:
Calibration measurements with gamma rays



VF Hess in his lab in 1915

In 1910 Hess became an assistant at the just-founded Radium Institute of the Imperial Academy of Sciences in Vienna. He performed absorption measurements in air with the strongest gamma source available at the institute and experimentally confirmed the absorption coefficient predicted by Eve.

He improved the electrometer's construction and developed a calibration method for electrometers using gauge radium sources of different strengths. For calibrated detectors from the company Günther & Tegetmeyer (Braunschweig), the accuracy when measuring the strength of unknown sources was about 5 per mil; uncalibrated instruments achieved 3% accuracy.

1911

VF Hess:
First three balloon flights

In August and October of 1911, Hess performed three balloon flights reaching altitudes of 200m to 1000m and confirmed the findings of Wulf, Bergwitz and Gockel. To prepare for a new series of flights, Hess designed and ordered improved instruments, two for gamma-ray detection and one with thin detector walls to measure beta rays.



Victor F Hess in the balloon's basket sometime between 1911 and 1912

1912

VF Hess:
Six balloon flights from the Prater in Vienna at lower altitudes

Six new flights were financed by the Imperial Academy of Sciences and supported with balloons from the Royal Imperial Austrian Aeronautical Club in Vienna. Hess measured the ionisation mainly with two or three electrometers:

- 17 April, during an eclipse of the sun at 1900m–2750m of altitude
- 26–27 April, at night for six hours at 300m–350m of altitude
- 20–21 May, at night at 150m–340m of altitude
- 3–4 June, at night at 800m–1100m of altitude
- 19 June, in the afternoon at 850m–950m of altitude
- 28 June, at night at 280m–360m of altitude


With the hydrogen-filled balloon Bohemia, provided by the German Aero Club in Bohemia, Hess, together with W Hoffory and E Wolf, reached an altitude of 5350m and landed at noon in Bad Saarow/Pieskow in Brandenburg. All three detectors measured a strong increase in ionisation.



Seven flight routes of VF Hess in 1912

7 Aug 1912

VF Hess:
Seventh balloon flight, reaching an altitude of 5350m
Discovery of cosmic rays



Electrometer used by VF Hess in 1912

Physik. Zeitschr. XIII, 1912. Hess, Durchdringende Strahlung bei sieben Freiballonfahrten. 1089

Tabelle der Mittelwerte.

Mittlere Höhe über dem Erdboden in m	Beobachtete Strahlung in Ionen pro ccm und sec.			
	Apparat 1		Apparat 2	
	Q ₁	Q ₂	Q ₃ (reduziert)	Q ₄ (nicht reduziert)
0	16,5 (13)	11,8 (20)	19,4 (5)	29,7 (9)
200	15,4 (13)	11,2 (19)	19,7 (5)	28,2 (9)
500	15,5 (9)	10,4 (6)	18,8 (5)	27,7 (5)
1000	15,5 (3)	10,3 (4)	19,8 (5)	28,3 (5)
2000	15,0 (7)	10,1 (9)	22,2 (4)	28,7 (4)
3000	17,3 (1)	12,3 (1)	31,2 (1)	22,5 (1)
5000	19,8 (1)	14,5 (1)	35,2 (1)	31,8 (1)
4000–1500	34,4 (1)	27,2 (1)	—	—

Mean values of all measurements during the seven flights at different altitudes (the number of ionisation values in brackets)

VF Hess summarised the results of these seven flights as follows:

- At altitudes of less than 1000m, the results are in general agreement with previous measurements.
- A radiation of high penetration power hits the atmosphere from above, which cannot be caused by radioactive emanations.
- This radiation contributes to the total amount of observed ionisation at lower altitudes as well.
- Assuming gamma radiation, the sun is not the source of the extraterrestrial radiation.
- There is no difference between ionisation measured during the day and at night.



1913—1926

Confirmation, Refusal, Confirmation

1913—1914 **W Kolhörster:**
Confirmation of the
discovery of cosmic rays




Werner Kolhörster from the University of Halle (Germany) took the important step towards very high altitudes. In 1913 he performed three flights with a hydrogen balloon up to 6000m and confirmed the results of Hess. Kolhörster started his record flight at the end of June 1914 with improved electrometers, reaching an altitude of 9300m. The ionisation reached $80.4 \text{ ions cm}^{-3} \text{ s}^{-1}$, and its increase (see figure) was a perfect confirmation that the high-energy radiation had an extraterrestrial origin. He estimated the absorption coefficient of this penetrating radiation to be smaller than the value for gamma rays from radium D by a factor of at least five.

W Kolhörster (at back centre) in 1913

1914 **A Gockel, VF Hess and M Kofler:**
High-altitude
measurements in the Alps

Gockel demonstrated the hardness of the radiation with measurements 6m below the surface of Lake Constance, in the Alps at Jungfrauoch and on the Aletsch Glacier at 3400m and 2800m of altitude.

Hess and Kofler performed long-term studies of the ionisation, as dependent on time and weather conditions. Two electrometers were installed on the 2044m-high Obir in the Austrian Alps.



Ionisation measured by VF Hess and W Kolhörster, as dependent on altitude

1915 **E von Schweidler:**
First discussion of different
assumptions about the
origin of cosmic rays



E von Schweidler

For the Festschrift marking the 60th birthdays of Elster and Geitel, a first theoretical investigation of possible sources of the cosmic radiation was presented by von Schweidler (Vienna). Based on the existing knowledge of ionising radiation, he excluded the upper atmosphere, the moon, the planets, the sun and other fixed stars as sources of cosmic rays. Schweidler concluded that "the less extreme requirements prefer the hypothesis of radioactive substances distributed in outer space."

1918 **VF Hess, W Schmidt:**
Estimate of the distribution
of radioactive gases in the
atmosphere

The model developed by Hess and Schmidt described a distribution of radioactive emanations in the atmosphere in agreement with experimental data. As shown in the table (at right), only radium D can reach the upper atmosphere. All other radioactive emanations are concentrated near the Earth's surface because of their short lifetimes.

The altitude where the fraction of radioactive emanations is reduced to 50% compared to the Earth's surface was estimated for different radioactive products:

Radium emanations and short-lived decays:	~ 1200m
Radium D and decay products:	< 10000m
Thorium emanations and thorium A:	2m—3m
Thorium B and decay products:	100m—150m
Actinium emanations and actinium A:	0.5m—1m
Actinium B and decay products:	10m—20m

1923 **RA Millikan et al:**
Radiation measurements
in the atmosphere

RM Otis (Pasadena), a co-worker of Millikan, observed with ionisation measurements in aeroplanes up to 5400m of altitude a dependence similar to that found by Hess and Kolhörster. However, the increase was smaller. RA Millikan and IS Bowen used an unmanned sounding balloon. A very light electrometer registered the ionisation, temperature and barometric pressure automatically on a photographic film. They measured only one averaged ionisation value between 5km and 15km of altitude, which was smaller than expected from the European results by about a factor of four. Millikan concluded that there is no radiation of cosmic origin with the absorption coefficient estimated by Hess and Kolhörster.



RA Millikan and GH Cameron in 1925

1926 **RA Millikan et al:**
Disproof of European
results, "rediscovery" of
cosmic rays


From measurements in snow-fed lakes at high altitudes, Millikan and Cameron deduced an absorption coefficient for the penetrating radiation in water supposedly smaller than found in Europe. Claiming the discovery of cosmic rays, Millikan caused strong reactions by Hess, Kolhörster and others.

1926—1933 Investigation of Properties

1926 L Myssowski, L Tuwim:
Barometric effect


The dependence of the ionisation rate on barometric air pressure was discovered in a three-week measurement. An electrometer was installed 1 m below the surface of the Neva River in Leningrad (now St Petersburg). An explanation for the observed increase in the rate with decreasing air pressure was given many years later when muons and pions were discovered in cosmic particle interactions.

The dependence of ionisation on air pressure (from Myssowski, Tuwim, *Physikalische Zeitschrift*, 39, 146 1926)



1927—1929 D Skobeltsyn:
First cosmic ray tracks in the cloud chamber


D Skobeltsyn in his laboratory in Leningrad



In 1927 Skobeltsyn investigated beta rays in a cloud chamber operating in a magnetic field. By chance, he observed in two pictures straight tracks which he interpreted as being due to high-energy cosmic rays. In a dedicated experiment, he found 36 tracks in 600 photographs. This was the first visual proof for the existence of charged secondaries produced by cosmic rays.

1927—1937 J Clay, AH Compton et al:
Latitude effect

A possible dependence of the cosmic ray rate on latitude was predicted by Kolhörster in 1919. Sailing in 1926 from Amsterdam to Indonesia, J Clay used an electrometer to measure the expected decrease of the ionisation near the equator. Through a worldwide measurement campaign initiated by Compton in the 1930s, the effect was established. This was the definitive proof that a part of the primary radiation consists of charged particles.





The relative cosmic particle intensity as a function of geomagnetic latitude

1928 H Geiger, W Müller:
New cosmic particle detector

In 1928 Geiger and Müller announced the development of a new detector. The counter consists of a metal tube filled with gas, and an isolated wire at a positive high voltage in the tube centre. If a particle crosses the tube, the gas becomes ionised and the electrons move to the wire, yielding a measurable signal.

1928 W Bothe, W Kolhörster:
Experiment with Geiger-Müller counters in coincidence




Absorber experiment with Geiger-Müller counters C1 and C2 in coincidence

Bothe and Kolhörster designed a trend-setting experiment to measure the absorption of cosmic rays with two Geiger-Müller counters in coincidence. The passage of a cosmic ray through both counters could be observed using electrometers. The absorption measurements were taken with and without a gold block between the counters. Coincidences could only be caused by particles, not by gamma rays. This was additional proof that at least a part of the secondary cosmic radiation consists of corpuscular particles.

1928 W Bothe: Invention of the coincidence circuit

Shortly after the absorption experiment, Bothe invented the electronic coincidence circuit. The concept was improved many times and is still an essential element in particle and astroparticle experiments.

Electronic coincidence circuit. There is a signal at C3 only if counters Z and Z' are crossed by the same particle.



1929 W Pauli:
Prediction of the neutrino


The neutrino was introduced by Pauli as a neutral particle to save the momentum conservation of the beta decay, which has to be a three-body decay. He assumed that it would never be detectable.

1929—1933 J Joly:
Biological impact

J Joly from the University of Dublin was probably the first to discuss possible biological impacts of cosmic radiation on the variation of species due to the interaction with chromosomes or on the incidence of cancer.

1930—1937 C Störmer, G Lemaitre, MS Vallarta:
Trajectories in the Earth's magnetic field

A visionary explanation for the appearance of the aurora borealis was given in 1898 by K Birkeland. Electrons emitted by solar flares are guided by the Earth's magnetic field to the polar areas and excite the molecules of the atmosphere. Based on this theory, Störmer (1930) and Lemaitre and Vallarta (1934) calculated the trajectories of cosmic particles.




The relative cosmic particle intensity as a function of geomagnetic latitude for minimum energies of electrons and protons

1931—1934 B Rossi, T Johnson, L Alvarez, AH Compton:
East-west effect



B Rossi (middle) in Eritrea



Geiger-Müller counter telescope


B Rossi's prediction of an east-west effect based on Störmer's calculations of cosmic particle trajectories in the Earth's magnetic field. Particles with a positive charge should enter the atmosphere from the west, those with a negative charge from the east. The effect was first demonstrated by Johnson and independently by Alvarez and Compton in 1932 at high altitude and low latitude in Mexico. Rossi confirmed the results in 1933 with his measurements in Eritrea.



1933—1947 Birth of Particle Physics


1932 **CD Anderson:**
Discovery of the positron

In 1931, with a cloud chamber operating in a strong magnetic field, Anderson observed cosmic ray tracks with negative and positive charges, which were interpreted as electrons and protons. Since many positive tracks had the same ionisation as the electrons, Anderson introduced a 6mm-thick lead plate into the chamber. In photographs from 1932, he found tracks with the ionisation and track length observed for electrons, but with a positive charge. This anti-electron (positron) had been predicted two years earlier by PAM Dirac.



A positron with an energy of 63MeV entering the lead plate from below and leaving the plate with an energy of 23MeV. For a proton, the track length would be ten times shorter.

1933—1935 **B Rossi, PMS Blackett, G Occhialini:**
Particle showers



Cloud chamber photograph of a particle shower with about 16 tracks. The divergence of the tracks points to an interaction in the magnet coil.


Rossi performed measurements with three Geiger-Müller counters in coincidence with and without lead shielding on top. The coincidence rate increased with the shielding, even though the opposite had been expected. The explanation was the shower production by an incoming cosmic particle. Blackett and Occhialini demonstrated the shower production visually with cloud chamber photographs.

1934 **W Baade, F Zwicky:**
Supernovae as possible sources of cosmic rays

By investigating photographic plates taken over the past 30 years, about 13 short flaring, extremely bright objects were identified. Zwicky and Baade called them supernovae. Based on the estimated energy release, they concluded that supernovae are sources of cosmic rays. This hypothesis is still valid, but not completely confirmed.


1935 **H Yukawa:**
Prediction of the pion

Yukawa formulated a theory to explain the dense packing of protons and neutrons in the nucleus of an atom. The short-ranged field needed a carrier with a mass inversely proportional to the range. He estimated a particle mass of about 100MeV and predicted that these particles could be produced in cosmic particle interactions.



H Yukawa, 1949

1936 **SH Neddermeyer, CD Anderson:**
Discovery of the muon



Stereographic photograph of a cloud chamber exposure. A muon enters the chamber from above and comes to rest below.


In a cloud chamber exposure with a 1 cm-thick platinum plate in the centre, 6000 photographs were taken. Anderson and Neddermeyer found about 25 events where the energy loss in the platinum absorber was much smaller than measured for electrons or positrons. Since the mass should be between the electron and proton masses, they first called it the mesotron. For several years, it was assumed that this particle was the predicted Yukawa particle.

1937 **M Blau, H Wambacher:**
First cosmic ray nuclear interaction in a photo emulsion

The photo-emulsion technique was developed by M Blau. In 1937 a five-month exposure to cosmic particles was performed at Hess's Hafelekar cosmic ray station at an altitude of 2300m. The discovery of a so-called star was a breakthrough of this detection technique. A cosmic particle interacted with an atom of the emulsion, producing eight tracks.

1938 **P Auger:**
Extensive air showers

With two Geiger-Müller counters in coincidence, Auger and his colleagues, Maze and Robley, detected extensive air showers. They measured the rate at up to 300m of counter distance and estimated the energy of the primary cosmic particles to be about 10^{15} eV.




A "star" produced in a photo emulsion by a cosmic particle

1942 **I Lange, SE Forbush:**
Solar cosmic particles

In February 1942, a large solar flare appeared. Lange and Forbush measured an increase in the cosmic particle rate of about 15%. They concluded that this additional fraction is caused by charged particles emitted by the solar flare.

1947 **DH Perkins, GPS Occhialini, CF Powell:**
Discovery of the pion

The pion event identified by Perkins. Tracks B and C are protons; D is a tritium nucleus. The short track E is a recoil nucleus. The grain density and scattering of track A correspond to a particle with a mass of about 100MeV.



In 1938 Yukawa and Sakata predicted the lifetime of the Yukawa particle to be about 10^{-8} seconds, which was 100 times shorter than the measured lifetime of the muon. The problem was solved with the discovery of the pion in photographic emulsions in 1947. Perkins found one event, and two months later Occhialini and Powell identified 25 pion interactions. In Britain, the emulsion technique was improved by Powell, Perkins and others, in cooperation with the Ilford company.

1947—1959 Extensive Air Showers

1947 First International Cosmic Ray Conference in Krakow



The first edition of the biennial International Cosmic Ray Conference was held in Krakow, Poland. In the front row of the photo stand P Blackett, J Blaton, A Wheeler and W Heitler. Other prominent participants included P Auger, G Bernardini, J Clay, M Cosyns, L Janossy, L Leprince-Ringuet and CF Powell.

1947 D Skobeltsyn et al: Start of air shower experiments in the Pamirs

After WWII, a broad cosmic ray research programme began in the Soviet Union. At 3860m of altitude, an experiment using Geiger counters was installed over an area of 1000m in diameter. For the first time a reduction of random signals was achieved by forming coincidences of local counters, followed by the requirement of coincidences of counters at a greater distance.

1949 E Fermi: Model of cosmic particle acceleration

The basic idea behind Fermi's model was that particles reach a higher energy when they enter the front of a plasma cloud which is moving with the very high velocity v . Such plasma clouds are produced, for instance, in supernova explosions. But the model cannot explain acceleration to very high energies.





Illustration of Fermi's acceleration model

1953 W Galbraith and JV Jelley: First air Cherenkov counter



The availability of very sensitive photon detectors (photomultipliers, or PMTs) allowed for proving PMS Blackett's hypothesis that cosmic air showers produce Cherenkov light. The British physicists W Galbraith and JV Jelley built the first very simple air Cherenkov counter using a rubbish bin. Working on cloudless nights, a mirror on the bottom of the bin focussed the incoming light on a PMT in front of the mirror. But it would take 36 years before the Cherenkov telescope detected high-energy gamma rays from the Crab Nebula.

First air shower Cherenkov counter


1954—1961 First generation of extensive air shower (EAS) arrays

In Britain an EAS array of 91 Geiger-Müller counters was built by TE Cranshaw and W Galbraith, covering an area of about 0.6km². It operated from 1954 to 1957 and measured primary energies of up to 10¹⁷eV. At MIT, a group led by B Rossi developed a pioneering detector

type. The fast timing of scintillation counters and PMTs allowed for the determination of shower direction and core position. Also the analysis techniques were the basis of future EAS experiments. Other strong activities were taking place in Japan and the Soviet Union.


1957 Launch of Sputnik 2, with Geiger counter for cosmic particle detection

The era of space experiments started with Sputnik 2. With a Geiger-Müller counter built by Vernov's group (Moscow State University), the intensity of cosmic rays was measured from 3—9 Nov. A strong increase in radiation was measured at a latitude of 60°. Later, it would be interpreted as the outer radiation belt.



Geiger-Müller counter installed in Sputnik 2

1958 J Van Allen: Discovery of belts of radiation




Van Allen discovered with Geiger counters installed on the Explorer 1 and Pioneer satellites that the Earth is surrounded by "clouds" of cosmic particles which are trapped by the Earth's magnetic field.

Van Allen belts surrounding the Earth

1958 Launch of Sputnik 3

This was the first mission with several novel scientific instruments which collected data over a period of two years. Cosmic rays were measured with a scintillation counter and their nuclear components with a Cherenkov counter.




Sputnik 3

1958 E Parker: Theory of the solar wind

The solar wind was originally discovered by L Biermann in 1951. Parker was the first to develop a theory of how electrons and protons can escape from the surface of the sun to follow the field lines of the sun's magnetic field. The solar wind was first measured by the Luna 1 satellite on its way to the lunar fly-by.

1958 NA Porter: Prototype of a water Cherenkov detector



Not only was the air Cherenkov detector built in Britain, but so was the first water Cherenkov detector. NA Porter used a water-filled steel tank with a photomultiplier looking from the top into the water. Relativistic charged particles produce Cherenkov light in the water, which is measured by the PMT. The construction from 1958 is not so different from the design of modern water Cherenkov detectors at IceTop and the Pierre Auger Observatory.

1959 GV Kulikov/GB Khristiansen: Discovery of the "knee" of the cosmic ray spectrum

With data from the Pamirs EAS and an earlier experiment, Kulikov and Khristiansen demonstrated that there is a change in the slope of the distribution of the number of particles per shower. They interpreted this discovery to mean that particles with energies greater than 10¹⁶eV (corresponding to more than 10⁷ particles per shower) are probably of metagalactic origin, now known not to be true.



1959—1970 Discoveries

**1959—1964 Chudakov et al:
First stereo Cherenkov telescope**



A stereoscopic Cherenkov telescope system was installed on the Crimean Peninsula to improve the sensitivity to and the reconstruction of the shower as well as the direction of the primary high-energy gamma. Twelve telescopes, each consisting of a container with a mirror and one photomultiplier, were installed on rails to vary the stereoscopic view. The sensitivity was low, but had they measured over several years, they might have been able to see the Crab Nebula.

Air Cherenkov light stereo telescope array

**1960 MA Markov:
Proposal of high-energy cosmic neutrino detection**


Markov proposed installing arrays of photomultipliers in deep lakes or in the sea to search for rare neutrino interactions. In about two-thirds of the interactions, a muon is produced which carries most of the energy and direction of the neutrino. Relativistic muons produce in water Cherenkov light which is detected by PMTs.

1960—1963 Volcano Ranch EAS experiment




An array of 19 scintillation detectors (3.26m² each) operated in the configuration at right (black dots in the schematic drawing) for three years at 1770m of altitude in New Mexico. It was the first experiment to measure air showers with energies greater than 10²⁰eV. The cosmic particle with the highest energy of 1.4 x 10²⁰eV (out of 5 x 10¹⁰ particles in the shower) was measured in February 1962. The experiment gave the earliest hint of a flattening of the energy spectrum at around 10²⁰eV.

Left: J Linsley searching for snakes in a detector station; right: hexagon structure of the array with the position of the UHE event



**1965 A Penzias and R Wilson:
Discovery of the cosmic microwave background radiation (CMB)**

The CMB radiation was detected by chance as Penzias and Wilson measured the background noise temperature of the Bell Labs horn-reflector antenna between July 1964 and April 1965. They found an excess temperature of 3.5°K which was isotropic, unpolarised and independent of seasonal variations. Dicke, Peebles, Roll and Wilkinson interpreted this effect as the cosmic microwave background radiation. The CMB is a relic of the early universe, which was a hot, dense plasma of photons and elementary particles. At that time, the temperature was about 3000°K, which cooled down due to the expansion of the universe.




Horn-reflector antenna

**1965 Kolar Gold Field and Case-Witwatersrand-Irvine:
Detection of atmospheric neutrinos**

The first atmospheric neutrinos, decay products of pions or kaons produced in interactions of cosmic particles in the atmosphere, were detected by two groups. The Bombay-Osaka-Durham collaboration operated a visual detector in the Kolar Gold Field of India at a depth of 7500m water equivalent. The Case-Witwatersrand-Irvine group worked in a South African gold mine at a depth of 8800m water equivalent. To reduce background muons, the detectors were set up to be triggered by horizontal muons.

**1965—1968 Launches of Proton 1—4:
Cosmic ray spectrum at 10¹¹—10¹⁵eV**



Ionisation calorimeter

For the first time an ionisation calorimeter built by Moscow State University was installed in satellites. This allowed a direct measurement of cosmic particle spectra in the energy range of 10¹¹eV to 10¹⁵eV below the “knee”.

**1966 K Greisen and G Zatsepin, V Kuzmin:
GZK cutoff energy for cosmic particles**

The prediction that the energy of cosmic particles from distant sources is limited to 5 x 10¹⁹eV was made independently by Greisen and by Zatsepin and Kuzmin. Cosmic particles with higher energies interact with photons of the cosmic microwave background radiation before they reach the Earth.

**1967—1987 Haverah Park:
EAS cosmic particle array**

To study EAS showers with energies from 10¹⁵eV to 10²⁰eV, a 12km² array of water Cherenkov counters was built and operated by the University of Leeds. For the first time, water Cherenkov counters were used. Together with Akeno in Japan, they found the “ankle” of the cosmic ray spectrum. The event with the highest energy was measured at 8.28 x 10¹⁹eV.

**1968 Whipple collaboration:
First 10m diameter Cherenkov telescope**




Whipple telescope

The 10m mirror of the Whipple telescope was a new level of quality in the design of air Cherenkov detectors which is still state of the art. In the first years of operation, a single PMT measured the Cherenkov light.

**1970 Yakutsk:
EAS array in Siberia**

Data-taking started in 1970 with a prototype array of 13 scintillation counter stations. The array was extended in the following years. In addition to scintillation counters, air Cherenkov and muon detectors were added. At 18km² it was at the time the largest, most complex array to measure the energy spectrum, the direction of primary particles and their mass composition in the energy range of 10¹⁷eV to 10²⁰eV.



Scintillation and air Cherenkov counter station



1971—1995 New Technologies


1973—1991 **Fly's Eye:**
First fluorescence detector array

Fluorescence light is produced when cosmic particle showers excite air molecules. The pioneering work was done by a group under K Greisen in the 1960s. Initial prototype detectors measuring air fluorescence light were built and tested by the University of Utah at the beginning of the 1970s. The Fly's Eye detector array was located in

the desert of Utah at an altitude of about 1370m. With 67 detector units, each consisting of a container with a 1.5m mirror and a light collection system, the array was able to register on moonless nights fluorescence light over an area of about 1000km². In 1991 an event was detected with $(3.2 \pm 0.9) \times 10^{20}$ eV, the highest energy ever measured.

1978—1996 **DUMAND project:**
Deep Underwater Muon and Neutrino Detector


Discussions about building DUMAND began in 1973. According to the 1978 proposal, one cubic kilometre of ocean was to be instrumented at a depth of 5km near the coast of Hawaii by about 23000 photomultiplier detectors. Neutrinos produce a muon in two-thirds of their very rare interactions. Relativistic muons produce in water Cherenkov light which is measured by PMTs. The arrival time allows for reconstructing the muon track. To disentangle the large amount of high-energy atmospheric muons from the muons produced in the few neutrino interactions, one looks for upward-going muons, as only neutrinos can traverse the Earth. The giant project failed for several reasons, but the almost 25 years of R&D were a very helpful basis for subsequent projects.



Muon detection through Cherenkov light

1984 **Baikal neutrino project:**
First stationary string


Lake Baikal in Siberia, with its clean water and stable ice in wintertime, was chosen for the installation of a water Cherenkov neutrino detector. The first string of PMT detectors, deployed at a depth of 1200m in 1984, measured atmospheric muons. Construction of the present NT-200 detector started at the end of the 1980s. An array of eight strings with 24 floors and 48 PMTs per string was planned. After several installation steps, NT-200 was completed in 1998. One of the first neutrino events with an upward-moving muon was detected in 1996.



Baikal NT-200 detector


1987 **Supernova 1987A:**
First detection of low-energy neutrinos

The neutrino window on the universe was opened by chance on 23 February 1987, when a supernova explosion in the Small Magellanic Cloud was discovered by optical telescopes. At the time several underground detectors were looking for proton decays in large volumes of water and in liquid scintillators. The data showed that three detectors observed an excess of events at the same time: the Japanese Kamiokande II detected 11 events within 13 seconds, the US IMB eight events within 6 seconds and the Soviet Union Baksan five events within 9 seconds from SN1987A.




Photomultipliers on the inner wall of Kamiokande II

1989 **Whipple collaboration:**
Gamma rays from the Crab Nebula



Crab Nebula

A telescope camera with 37 PMTs and an efficient analysis algorithm allowed a considerable reduction of hadronic background events. In 1989, more than 20 years after the Whipple telescope started to look for Cherenkov light flashes, the discovery of gamma rays from the Crab Nebula opened a new window into the sky: high-energy gamma astronomy.



1993—2009 **AMANDA neutrino detector**

The first string of photomultiplier modules was installed in the ice at the South Pole in the austral summer of 1993—1994. The AMANDA detector collected data in the final configuration of 19 strings from 2000—2009. About 1000 muon-neutrino interactions per year were measured. However, the neutrinos were produced in pion decays in the atmosphere of the northern sky. No significant neutrino signal from a galactic or extragalactic point source was found.


Schematic view of the 19-string AMANDA detector mostly installed at a depth of 1400m—2000m

1993 **AGASA:**
Highest-energy event


The Akeno Giant Air Shower Array (AGASA) operated in Japan from 1990—2004. It consisted of 111 scintillation and 27 muon detectors distributed over an area of 100km². The event with the highest energy of 2×10^{20} eV was measured in December 1993. The particle shower was distributed over an area of 6x6km². In total, 11 events with an energy greater than 10^{20} eV were detected; this did not show the decrease expected for the GZK effect observed in other experiments.

1995 **Start of SOHO:**
Solar and Heliospheric Observatory

SOHO is a joint project of NASA and ESA to study the sun and the solar wind. Originally designed to operate for two years, it still provides important information about the structure of sunspots and the temperature profile and gas flow in the corona over a complete sun cycle of 11 years. SOHO is also a very important detector for space weather, providing alerts in the case of sun bursts directed at the Earth.




Solar burst seen by SOHO



SOHO satellite

1995 **Detection of antiprotons**
in primary cosmic rays

In the 1970s, NASA began the exploration of primary cosmic rays with balloon experiments at very high altitudes. Particle detectors weighing more than 1000kg measured particle energies and the composition of the primaries, and searched for antimatter. Three experiments launched in the early 1990s, IMAX, BESS and CAPRICE, published the discovery of antiprotons.



IMAX magnet spectrometer. TOF - time of flight, C1/C3 - Cherenkov detectors, S1/S2 - scintillators, MWPC - multi-wire proportional chamber, DC1/DC2 - drift chambers.



1996—2012 Gamma-Ray Astronomy

1996—2010 **KASCADE & KASCADE-Grande**



KASCADE detector array

With the Karlsruhe Shower Core and Array DEtector, a new generation of modern shower arrays came into operation. Some 252 stations, each consisting of four electron/gamma liquid scintillation counters and a muon counter, were arranged in an array of $200 \times 200 \text{ m}^2$. In the centre a hadron and muon detector system with an active area of 320 m^2 was installed. KASCADE was extended in 2003 by 37 scintillation counter stations to become KASCADE-Grande. This allowed the study of the composition of primary cosmic particles from 10^{15} eV to 10^{18} eV , covering the “knee” and the so-called “second knee”.

1997—2006 **HiRes**

The High Resolution air fluorescence experiment, known as HiRes, replaced the first-generation Fly’s Eye detectors in Utah. With smaller photomultipliers and an improved data acquisition system, a higher sensitivity could be reached in order to determine the energy, direction and chemical composition of the highest-energy cosmic rays. At energies greater than $5 \times 10^{19} \text{ eV}$, the decrease in the measured particle flux is in agreement with the expectations of the GZK effect.

2002 **Third generation of Cherenkov telescopes**



H.E.S.S.: 4 telescopes in 2003, 5 in 2012




MAGIC: 1 telescope in 2004, 2 in 2009



VERITAS: 2 telescopes in 2006, 4 in 2008


The success story of high-energy gamma astronomy began with the operation of the new generation of air Cherenkov telescopes: H.E.S.S. in Namibia, MAGIC in La Palma and VERITAS in Arizona.

2004 **IceCube & IceTop**



Digital optical module

From 2004—2010, the giant IceCube Neutrino Observatory was built at the South Pole. Some 86 strings with digital optical modules were installed to survey one cubic kilometre of ice to depths of 1450m to 2450m. Every hour about 10 up-going muons from neutrino interactions near the detector and about 7 million down-going atmospheric muons are reconstructed by measuring the Cherenkov light. In addition, 81 pairs of IceTop tanks are installed on the surface to investigate air showers in the “knee” energy region. In the data analysed so far, no neutrinos have been observed from point sources. All the neutrinos measured since 2005 were produced in the northern atmosphere.



IceCube detector

2004 **Pierre Auger Observatory**




The largest extensive air shower array was constructed as the first true hybrid observatory. It comprises 1600 water Cherenkov detectors distributed on a grid of 1.5 km over an area of 3000 km^2 , and 27 air fluorescence telescopes erected at four stations on the periphery of the observatory. Important results include the exploration of the ultra-high-energy area with a cut-off as expected from the GZK effect; the particle composition; and a first hint of the extragalactic origin of the highest-energy cosmic particles.

One of the air fluorescence light detector stations on the hill and a water Cherenkov tank on the right

2006 **PAMELA mission**

The Italian-Russian satellite was launched with the goal of antimatter-matter exploration in primary cosmic rays. PAMELA measured an unexpected increase in the positron fraction at energies between 10^{10} eV and 10^{11} eV which was later established by Fermi LAT at even higher energies. Further investigations will answer whether this effect opens a window on a “new physics”.




Schematic view of the PAMELA detector

2008 **Telescope Array**

The Telescope Array project in Utah combines the successful tradition of Fly’s Eye and HiRes air fluorescence shower detection with a ground array of scintillator counters distributed over an area of about 500 km^2 . The experiment is complemented by a low-energy extension. Both the particle flux and composition can be studied for primary particle energies above $3 \times 10^{19} \text{ eV}$.

2008 **Fermi Large Area Telescope and Burst Monitor**

Fermi LAT all-sky view




The Fermi satellite carries two experiments, the Large Area Telescope (LAT) to measure high-energy gamma rays up to energies of 10^{11} eV and the GLAST Monitor for the detection of gamma-ray bursts. The high sensitivity of both instruments allowed the detection of many new galactic and extragalactic gamma-ray sources (with LAT) and gamma-ray bursts (with GLAST). With the LAT electromagnetic calorimeter, the electron flux was measured up to energies of 10^{12} eV . For positrons, the energy range from $2 \times 10^{10} \text{ eV}$ to $2 \times 10^{11} \text{ eV}$ was investigated and an increase in the positron fraction observed.

2010 **Evolution of the gamma-ray sky**

The progress of very high-energy gamma ray astronomy with ground-based air Cherenkov telescopes over the last two decades can be seen in the sky plots of detected sources. With the second-generation Whipple telescope, one source, the Crab Nebula, was observed until 1990.

Ten years later, Whipple and HEGRA had seen nine sources. And from 2003—2010, the third-generation telescopes H.E.S.S., MAGIC and VERITAS detected about 110 gamma sources.



2011 **Launch of the Antimatter-Matter Spectrometer**

The AMS cosmic particle spectrometer installed at the International Space Station was designed to search for cosmic particles and antiparticles up to energies of 10^{12} eV . A further goal is the indirect search for dark matter. To search for possible anti-matter concentrations in the universe predicted by some theories, the sensitivity for measuring the anti-helium/helium fraction was improved by a factor of 1000.

