

Physics at the LHC

Lecture 13: New Physics (non-SUSY) scenarios at the LHC

Klaus Mönig, Sven Moch


(klaus.moenig@desy.de, sven-olaf.moch@desy.de)



Wintersemester 2010/2011

Addendum SUSY

- Meanwhile ATLAS and CMS have made their 1st SUSY results public, CMS in the jets+MET, ATLAS in the jets+MET+1-lepton channel
- This excludes masses up to $m_{\tilde{q}} < 700 \text{ GeV}$ for a specific choice of parameters
- More generic limits will come later



Reasons for new physics

The hierarchy problem: How to stabilise the Higgs mass with $M_{\text{pl}} \sim 10^{17} m_{\text{H}}$

- **SUSY**: contributions from particles and superpartners cancel
- **Extra dimensions (ADD type)**: in reality $M_{\text{pl}} \sim m_{\text{H}}$ but gravity propagates in $3+n$ dimensions
- **Extra dimensions (RS type)**: a 4th dimension with a “warped” geometry separates the Planck brane with $M_{\text{pl}} \sim m_{\text{H}}$ and our brane (TeV brane)
- **Little Higgs**: The loops are cancelled by new SM-like particles at one loop, new physics in the 10 TeV range
- **Technicolour**: the Higgs mechanism is mimicked by new strong interactions at the TeV scale

What is the origin of dark matter?

- We expect a particle with $0.1 - 1$ TeV mass, neutral, weakly interacting
- Many models contain such a particle which can be made stable with a special parity

Where does the baryon-antibaryon asymmetry come from?

- We can have an additional source of CP violation
- With a more complicated neutrino sector one can first generate a lepton-antilepton asymmetry that is then transferred to the baryon sector
- This requires lepton/baryon number violating interactions

Strategies to find new physics

- Find the Higgs and measure its properties
- Look for particles that are predicted by specific (classes of) models
- Look for generic signals like missing E_T from dark matter production
- Look for an extended gauge sector (Z' , W') predicted by many models
- Measure properties and interactions of gauge bosons, top quarks etc.

Search for new gauge bosons



- Many models contain an enhanced gauge group
- The new interactions result in new gauge bosons
- Most models contain a neutral Z'
- Many models also contain a charged W'
- At LHC visible via $q\bar{q} \rightarrow \ell^+ \ell^-$ and $q\bar{q}' \rightarrow \ell \nu$
- ⇒ Z' cross section $\propto (g_V(q)^2 + g_A(q)^2) (g_V(\ell)^2 + g_A(\ell)^2) / \Gamma$
(interference with Drell Yan plays a minor role)
- The sensitivity gets an additional $1/\Gamma$ factor from the background scaling
- ⇒ Relatively weak model dependence

Z' searches

Event selection:



- two high energy, opposite sign, same flavour leptons
- if needed some isolation criteria

TEVATRON limits $m(Z') \gtrsim 1 \text{ TeV}$ depending on model



At LHC $m(Z') \gtrsim 3 \text{ TeV}$ with relatively low luminosity


Further improvement difficult due to steeply falling PDFs




At 8 TeV Z' discovery for the most favourable case (SSM) up to 2 — 2.5 TeV is possible.

How can one identify the model?

- The total cross section is proportional to $\left(g_{V,q}^2 + g_{A,q}^2\right) \left(g_{V,\ell}^2 + g_{A,\ell}^2\right) / \Gamma$
 - The total width can be fitted from the data and is $\propto \sum_i \left(g_{V,i}^2 + g_{A,i}^2\right)$
 - To get information on the couplings from the width all decay modes must be known
 - The proton contains two valence up-quarks and only one valence down-quark
 - $\Rightarrow$ at high x up-quarks dominate
 - $\Rightarrow$ at high Z' rapidity more Z' originate from up-quarks
- $\Rightarrow$ The Z' rapidity distribution contains information in the up/down-quark coupling ratio





- The forward backward asymmetry depends on the ratio of the vector and axial vector coupling
- At high mass usually the boost direction determines the direction of the quark
- This gives some distinction between the models if the mass is not too high



W' searches

- Only one lepton and missing E_T from neutrino
- Longitudinal ν -momentum unknown $\Rightarrow$ only m_T can be calculated
- In general W' cross section larger than Z' cross section
- Event selection: one (isolated) high energy lepton, missing E_T and some jet veto
- Reach $\sim 0.5 - 1$ TeV higher than Z'



Models with extra dimensions

Several models contain extra dimensions

- Large extra dimensions (ADD):
 - several (2-7) extra dimensions
 - extra dimensions are large (μm – nm)
 - only gravitation can live in the bulk
- Universal extra dimensions:
 - all fields live in the bulk
 - this limits the size of the extra dimensions to several hundred GeV
- Randall Sundrum models
 - one extra dimension with warped geometry
 - gravity is located on different brane than TeV physics
 - only gravity or all fields can live in the bulk
- String theories
 - in general no visible signal since extra dimensions on Planck scale

ADD type extra dimensions

- Only gravity lives in the bulk
- Size of the extra dimensions is $\mathcal{O}(\text{eV})$ or $\text{nm} - \mu\text{m}$
- This is also the spacing of the KK graviton resonances
- For LHC energies this is a continuous spectrum of resonances
- Physics interest:
 - In reality Planck mass is on TeV scale
 - Planck mass appears so large because gravity escapes into extra dimensions


$$M_{\text{pl}}^2 = M_{\text{D}}^{2+n} R^n \quad \left(\Rightarrow R \sim 10^{\frac{30}{n}-17} \left(\frac{1 \text{ TeV}}{M_{\text{D}}} \right)^{1+\frac{2}{n}} [\text{cm}] \right)$$

R : compactification radius of extra dimensions


- Experimentally get limit on R for assumed n
- This is turned into a limit on M_{D}

Visible processes:

Virtual Graviton Exchange




Direct Graviton Production



- The cross section for a single KK graviton is negligibly small
- However due to the large number of excitations the total effect has the size of an electroweak cross section


Limits from the Tevatron

- At the Tevatron analyses are done with jets + missing E_t and Drell-Yan type events
- The analyses give limits around 1 TeV almost independent of the number of extra dimensions
- LEP has analysed the data with one photon and missing energy
- The LEP limits depend stronger on the number of dimensions




Expected effects

Virtual graviton exchange:
Expect broad enhancement of
Drell Yan production




Graviton radiation:
Jet events with large missing energy





Both effects are sensitive to $M_D \lesssim 6 - 9 \text{ TeV}$

Black holes at the LHC


- Black hole production is possible, when the centre of mass energy gets into the region of the Planck mass
- This would be fulfilled for ADD models
- Schwarzschild-Radius
$$R_S = \frac{2GM_{BH}}{c^2}$$
- Cross section $\sigma \sim \pi R_S^2 \sim 100 \text{ pb}$
- Black holes decay by Hawking radiation with
$$\tau_{BH} \sim 10^{-27} - 10^{-25} \text{ s}$$
- Decay is thermal: high multiplicity, symmetric, democratic in particle species ($\Rightarrow$ leptons, neutrinos)





Black holes are easy to separate from background by lepton number, $\sum p_T$ and by event shape variables that are sensitive to the isotropy of the decay



From other new physics like SUSY they can be separated e.g. with missing E_T





- Black holes can be found up to around 10 TeV with almost no luminosity
- The mass can be reconstructed from the visible decay products
- Some information on the number of extra dimensions can be obtained from other variables like decay-multiplicity



Black holes at 7 TeV


- Limits on black holes exist around 4 TeV
- This assumes, however, optimistically large cross sections



Randall Sundrum models



- Two branes separated in a 5th dimension with an deSitter geometry
- Mass scale on SM brane exponentially suppressed w.r.t. Planck brane
 $ds^2 = e^{-2ky} \eta_{\mu\nu} dx^\mu dx^\nu + dy^2 \quad y = \phi r_c$
- Scale $\Lambda = M_{Pl} e^{-kr_c \pi} \sim 1 \text{ TeV}$
- Equally spaced KK resonances with mass $m_1 = x_1 k / M_{Pl} \Lambda$
 $0.01 < k/M_{Pl} < 0.1$

- Original version only gravitation in the bulk
- KK-graviton decays symmetric in flavours
- Graviton has spin 2





Search for RS-Gravitons

- Most efficient search is in leptonic decay channel
- In this channel the search is identical to the Z' search
- At low luminosity 1-2 TeV, at high luminosity 2-3 TeV reach are possible
- This excludes the entire region allowed by the model




How to identify that this is a KK graviton and not a Z'

- Gravitons have spin two resulting in a different decay angle distribution
- For the decay angle distribution it is important that the KK-Graviton can also be produced by gluons
- In most of the parameter space spin 1 can be excluded



RS with matter in the bulk


- New versions of the RS model allow matter in the bulk
- A new parity can produce dark matter
- The matter fields are located in different positions in the bulk generating the large mass differences in the SM
- This causes the KK resonances to decay dominantly into heavy fermions
- First studies show large effects from the KK graviton, but only very small effects from the γ and Z excitation




Universal extra dimensions

- If all fields live in the bulk the compactification scale must be at least a few hundred GeV
- A KK parity can be defined that forbids even-odd transitions of KK resonances and allows even-even and odd-odd transitions only on loop level
- If the higher resonances are not seen the model is very SUSY like

M. Peskin, Victoria




- Most efficient search: jets plus missing E_T
- This allows to find UED scenarios up to 2.7 TeV




Search for higher resonances

- For one extra dimension the next resonance has twice the mass of the first one
- For two extra dimensions the $(1,1)$ resonance has $\sqrt{2}$ times the mass of the first resonance
- The $(1,1)$ resonances should have a large branching ratio to $t\bar{t}$
- Unfortunately the cross section is still low
- According to present studies the LHC can find these resonances only up to around 1.5 TeV




Strong Electroweak Symmetry Breaking

- If no Higgs exists electroweak interactions become strong at high energy and e.g. WW scattering violates unitarity at $\sqrt{s_{WW}} \sim 1.2 \text{ TeV}$.
- ⇒ expect new effects at this energy
- Typical models invoke a new strong interaction at the TeV scale (Technicolour)
- The Goldstone-bosons (Pions) of the new theory become the longitudinal degrees of freedom of the vector-bosons
- Warning: simple copy of QCD is excluded by LEP/SLD precision data



Systematic treatment





Effective Lagrangian for VV scattering:

$$\mathcal{L}_4 = \alpha_4 \left[\frac{g^4}{2} \left[(W_\mu^+ W^{-\mu})^2 + (W_\mu^+ W^{+\mu})(W_\nu^- W^{-\nu}) \right] \right. \\ \left. + \frac{g^4}{c_w^2} (W_\mu^+ Z^\mu)(W_\nu^- Z^\nu) + \frac{g^4}{4c_w^4} (Z_\mu Z^\mu)^2 \right]$$

$$\mathcal{L}_5 = \alpha_5 \left[g^4 (W_\mu^+ W^{-\mu})^2 + \frac{g^4}{c_w^2} (W_\mu^+ W^{-\mu})(Z_\nu Z^\nu) \right. \\ \left. + \frac{g^4}{4c_w^4} (Z_\mu Z^\mu)^2 \right]$$


(assuming custodial symmetry, C, P conservation)

- Effective theory valid up to $\Lambda = 4\pi v$
- Expect $|\alpha_i| \lesssim v/\Lambda = 1/4\pi$
- Need unitarisation procedure, e.g. Pade unitarisation (works well in meson physics)
- Most likely get resonances (like ρ and ω in QCD)
- However also models without resonances are possible



Event selection:

- At least one W,Z decays leptonically
- One W or Z can decay into one or two jets
- For ZZ one Z can decay into neutrinos
- Forward jet tagging and central jet veto similar to Higgs fusion channel
- Sensitivity up to around 1 TeV no results for no-resonance case yet




Little Higgs models

- In the SM the Higgs-mass receives large loop corrections
 - from the top loop $\delta m_h^2 = \frac{3}{8\pi^2} \lambda_t^2 \Lambda^2 \sim (2 \text{ TeV})^2$ ($\Lambda = 10 \text{ TeV}$)
 - from the W/Z loops $\delta m_h^2 \sim \alpha_w \Lambda^2 \sim -(750 \text{ GeV})^2$
 - from the Higgs loop $\delta m_h^2 \sim \frac{\lambda}{16\pi^2} \Lambda^2 \sim -(1.25 m_h)^2$
- The SM is embedded in a larger gauge group at $\Lambda_H = \mathcal{O}(10 \text{ TeV})$
- The Higgs is a pseudo-Goldstone boson of this breaking
- This protects the Higgs from one-loop corrections $\propto \Lambda_H^2$
- Technically this is done by new particles of same spin and $\mathcal{O}(1 \text{ TeV})$ mass:
 - An extended Higgs sector
 - New gauge bosons $A_H Z_H, W_H^\pm$ with mass $\lesssim 6 \text{ TeV} \left(\frac{m_H}{200 \text{ GeV}}\right)^2$
 - A new top-like quark T with mass $\lesssim 2 \text{ TeV} \left(\frac{m_H}{200 \text{ GeV}}\right)^2$

Search for the new gauge bosons



Discovery channel: leptonic decays

- Search equal to Z' , W' searches already shown
- Cross section in general very large depending on a mixing angle θ
- Discovery up to ~ 6 TeV depending on mixing angle




How to identify the little Higgs model?

- Decays to heavy quarks are predicted by the model
- Measurement of them helps to identify the model
- Can be seen up to 3 TeV






- Very important W_H , Z_H decay to Higgses which are needed to cancel the W and Z loops
- Can use $W_H \rightarrow WH \rightarrow \ell \nu b \bar{b}$, $Z_H \rightarrow ZH \rightarrow \ell \ell b \bar{b}$, $W_H, Z_H \rightarrow W, Z H \rightarrow jj \gamma \gamma$
- However decay only visible in limited parameter space




Search for the T-quark

- T and t mix with mixing parameter λ_1/λ_2
- Production via $qq, gg \rightarrow T\bar{T}$ or $qb \rightarrow q'T$, cross sections depend on λ_1/λ_2




- T decays dominantly in Zt , Wb and ht
- Search for Zt , Wb with leptonic W,Z decays

- Wb can be reconstructed with high efficiency but significant background
- Zt much cleaner but lower efficiency
- Mass reach 1-2 TeV depending on model parameters



Fermion substructure

- The fermions can have a substructure at a high scale
- In this case excitations of the fermions should exist
- Also the scattering cross section should be modified
- The interaction can be parametrised with a contact interaction





$$\mathcal{L}_{eff} = \frac{4\pi}{\Lambda^2} \sum_{i,k=L,R} \alpha^{ik} (\bar{q}_i \gamma^\mu q'_i) (\bar{f}_k \gamma^\mu f'_k)$$

$\Rightarrow$ cross section rises with s



- Λ : contact interaction scale. Must be $>$ few TeV from previous experiments

Excited fermions

- Excited fermions could exist at a scale around or smaller than Λ
- Single and pair production is possible
- Single production has a much larger cross section due to the smaller needed s'
- They decay into a fermion and a gauge boson (γ, W, Z)
- The mass reach is 4-7 TeV for 300 fb^{-1}




Already with 1/10th of last year's luminosity a limit around 1.5 TeV could be set.



Quark substructure


- Quark substructure will also be visible in the inclusive jet cross section at large p_T
- Effect gets large close to the compositeness scale



- Statistically no problem to get up to around $\Lambda = 40 \text{ TeV}$



However systematics are an issue:

PDFs uncertainties start to get relevant but are not the limiting factor





Limiting systematics: linearity of energy scale

- At lower energy the jet-energy scale can be measured very precisely from Z+jets events
- At higher energy the scale can be estimated from boot-strap methods e.g. using three jet events
- This introduces some non-linearity in the energy scale
- Since the QCD cross section falls so steeply this introduces a large uncertainty in the jet rate
- A 2% non linearity at 2 TeV limits the sensitivity to $\Lambda = 10 - 20$ TeV




Also hear 3.1 pb^{-1} where sufficient to set a limit around 3.5 TeV





Leptoquarks

- Grand unification models try to embed the SM gauge group into a larger group like $SU(5)$
- Such a group contains new interaction which are lepton- and baryon-number violating
- The corresponding gauge bosons have baryon and lepton number $\neq 0$ and decay into quark-lepton (leptoquarks)
- Some production channels:





Cross section

- For low masses the cross sections are very high
- However they fall steeply for higher masses
- In general the single LQ cross section is higher than the pair-production one



Event selection

- Two same flavour leptons with high p_t
- High E_T jets
- Large $\ell\ell$ mass to suppress Drell-Yan
- With these cuts a clean selection is possible
- Leptoquarks up to ~ 800 GeV can be discovered with very low luminosity
- For high luminosity only about 1.5 TeV are possible due to fast falling cross section



Model independent searches

Idea:

- Plot as many distributions as possible and compare with prediction
- New physics will show up in deviations of the data from the prediction

Problem:

- Impossible to understand all variables at hadron colliders with good precision
- All problems in detector understanding will appear as deviations in the plots (can try to fit some correction factors)
- With many variables expect some deviations from statistical fluctuations (can be corrected for)

Advantages:

- Model independent
- Don't overlook unexpected new physics

Disadvantages:

- Non-optimal cuts since one wants to stay model independent
- ⇒ signals may remain hidden under background

The CDF VISTA approach

- Plot as many distributions as possible
- Derive correction factors from a fit
- Correct significance for trial factor

Global result


CDF Run II Preliminary (2.0 fb⁻¹)
The calculation of σ accounts for the trials factor

Final State	Data	Background	σ	Final State	Data	Background	σ	Final State	Data	Background	σ
b $e^{\pm}\bar{p}$	690	817.7 $\pm$ 9.2	-2.7	2j $\bar{p}$ high- Σp_T	87	80.9 $\pm$ 6.8	0	j $\mu^{\pm}\mu^{\mp}\bar{p}$	32	32.2 $\pm$ 10.9	0
$\gamma\tau^{\pm}$	1371	1217.6 $\pm$ 13.3	+2.2	2j $\bar{p}$ low- Σp_T	114	79.5 $\pm$ 100.8	0	j $\mu^{\pm}\mu^{\mp}\gamma$	14	11.5 $\pm$ 2.6	0
$\mu^{\pm}\tau^{\pm}$	63	35.2 $\pm$ 2.8	+1.7	2j $\bar{p}\tau^{\pm}$	18	13.2 $\pm$ 2.2	0	j $\mu^{\pm}\mu^{\mp}$	4852	4271.2 $\pm$ 185.4	0
b2j $\bar{p}$ high- Σp_T	255	327.2 $\pm$ 8.9	-1.7	2j $\gamma\tau^{\pm}$	142	144.6 $\pm$ 5.7	0	j $\mu^{\pm}$	77689	76987.5 $\pm$ 930.2	0
2j $\tau^{\pm}$ low- Σp_T	574	670.3 $\pm$ 8.6	-1.5	2j $\gamma\bar{p}$	908	980.3 $\pm$ 63.7	0	$e^{\pm}4j\bar{p}$	903	830.6 $\pm$ 13.2	0
3j $\tau^{\pm}$ low- Σp_T	148	199.8 $\pm$ 5.2	-1.4	2j γ	71364	73021.4 $\pm$ 595.9	0	$e^{\pm}4j\gamma$	25	29.2 $\pm$ 3.6	0
$e^{\pm}\bar{p}\tau^{\pm}$	36	17.2 $\pm$ 1.7	+1.4	2j $\mu^{\pm}\tau^{\mp}$	16	19.3 $\pm$ 2.2	0	$e^{\pm}4j$	15750	16740.4 $\pm$ 390.5	0
2j $\tau^{\pm}\tau^{\mp}$	33	62.1 $\pm$ 4.3	-1.3	2j $\mu^{\pm}\bar{p}$	17927	18340.6 $\pm$ 201.9	0	$e^{\pm}3j\tau^{\mp}$	15	21.1 $\pm$ 2.2	0
$e^{\pm}j$	741710	764832 $\pm$ 6447.2	-1.3	2j $\mu^{\pm}\gamma\bar{p}$	31	27.7 $\pm$ 7.7	0	$e^{\pm}3j\bar{p}$	4054	4077.2 $\pm$ 63.6	0
j2 $\tau^{\pm}$	105	150.8 $\pm$ 6.3	-1.2	2j $\mu^{\pm}\gamma$	57	58.2 $\pm$ 13	0	$e^{\pm}3j\gamma$	108	79.3 $\pm$ 5	0
$e^{\pm}2j$	256946	249148 $\pm$ 2201.5	+1.2	2j $\mu^{\pm}\mu^{\mp}\bar{p}$	11	7.8 $\pm$ 2.7	0	$e^{\pm}3j$	60725	60409.3 $\pm$ 723.3	0
2bj low- Σp_T	279	352.5 $\pm$ 11.9	-1.1	2j $\mu^{\pm}\mu^{\mp}$	956	924.9 $\pm$ 61.2	0	$e^{\pm}2\gamma$	41	34.2 $\pm$ 2.6	0
j $\tau^{\pm}$ low- Σp_T	1385	1525.8 $\pm$ 15	-1.1	2j $\mu^{\pm}$	22461	23111.4 $\pm$ 366.6	0	$e^{\pm}2j\tau^{\pm}$	37	47.2 $\pm$ 2.2	0
2b2j low- Σp_T	108	153.5 $\pm$ 6.8	-1	2e $^{\pm}j$	14	13.8 $\pm$ 2.3	0	$e^{\pm}2j\tau^{\mp}$	109	95.9 $\pm$ 6.8	0
b $\mu^{\pm}\bar{p}$	528	613.5 $\pm$ 8.7	-0.9	2e $^{\pm}e^{\mp}$	20	17.5 $\pm$ 1.7	0	$e^{\pm}2j\bar{p}$	25725	25403.1 $\pm$ 209.4	0
$\mu^{\pm}\gamma\bar{p}$	523	611 $\pm$ 12.1	-0.8	2e $^{\pm}$	32	49.2 $\pm$ 3.4	0	$e^{\pm}2j\gamma\bar{p}$	30	31.8 $\pm$ 4.8	0
2b γ	108	70.5 $\pm$ 7.9	+0.1	2b high- Σp_T	666	689 $\pm$ 9.4	0	$e^{\pm}2j\gamma$	398	342.8 $\pm$ 15.7	0
8j	14	13.1 $\pm$ 4.4	0	2b low- Σp_T	323	313.2 $\pm$ 10.3	0	$e^{\pm}2j\mu^{\mp}\bar{p}$	22	14.8 $\pm$ 1.9	0
7j	103	97.8 $\pm$ 12.2	0	2b3j low- Σp_T	53	57.4 $\pm$ 6.5	0	$e^{\pm}2j\mu^{\mp}$	23	15.8 $\pm$ 2	0
6j	653	659.7 $\pm$ 37.3	0	2b2j high- Σp_T	718	803.3 $\pm$ 12.7	0	$e^{\pm}\tau^{\pm}$	437	387 $\pm$ 5.3	0
5j	3157	3178.7 $\pm$ 67.1	0	2b2j $\bar{p}$ high- Σp_T	15	21.8 $\pm$ 2.8	0	$e^{\pm}\tau^{\mp}$	1333	1266 $\pm$ 12.3	0
4j high- Σp_T	88546	89096.6 $\pm$ 935.2	0	2b2j γ	32	39.7 $\pm$ 6.2	0	$e^{\pm}\bar{p}\tau^{\mp}$	109	106.1 $\pm$ 2.7	0
4j low- Σp_T	14872	14809.6 $\pm$ 186.3	0	2b2j $\mu^{\pm}\bar{p}$	14	17.3 $\pm$ 1.9	0	$e^{\pm}\bar{p}$	960826	956579 $\pm$ 3077.7	0
4j2 γ	46	46.4 $\pm$ 3.9	0	2b2j $\mu^{\pm}$	22	21.8 $\pm$ 2	0	$e^{\pm}\gamma\bar{p}$	497	496.8 $\pm$ 10.3	0
4j $\tau^{\pm}$ high- Σp_T	29	26.6 $\pm$ 1.7	0	2b $\mu^{\pm}\bar{p}$	11	14.4 $\pm$ 2.1	0	$e^{\pm}\gamma$	3578	3589.9 $\pm$ 24.1	0
4j $\tau^{\pm}$ low- Σp_T	43	63.1 $\pm$ 3.3	0	2bj high- Σp_T	891	967.1 $\pm$ 13.2	0	$e^{\pm}\mu^{\pm}\bar{p}$	31	29.9 $\pm$ 1.6	0
4j $\bar{p}$ high- Σp_T	1064	1012 $\pm$ 62.9	0	2bj $\bar{p}$ high- Σp_T	25	31.3 $\pm$ 3.1	0	$e^{\pm}\mu^{\mp}\bar{p}$	109	99.4 $\pm$ 2.4	0
4j $\gamma\tau^{\pm}$	19	10.8 $\pm$ 2	0	2bj γ	71	54.5 $\pm$ 7.1	0	$e^{\pm}\mu^{\pm}$	45	28.5 $\pm$ 1.8	0
4j $\gamma\bar{p}$	62	104.2 $\pm$ 22.4	0	2bj $\mu^{\pm}\bar{p}$	12	10.7 $\pm$ 1.9	0	$e^{\pm}\mu^{\mp}$	350	313 $\pm$ 5.4	0
4j γ	7962	8271.2 $\pm$ 245.1	0	2be $^{\pm}2j\bar{p}$	30	27.3 $\pm$ 2.2	0	$e^{\pm}j2\gamma$	13	16.1 $\pm$ 3.9	0
4j $\mu^{\pm}\bar{p}$	574	590.5 $\pm$ 13.6	0	2be $^{\pm}2j$	72	66.5 $\pm$ 2.9	0	$e^{\pm}j\tau^{\mp}$	386	418 $\pm$ 18.9	0
4j $\mu^{\pm}\mu^{\mp}$	38	48.4 $\pm$ 6.2	0	2be $^{\pm}\bar{p}$	22	19.1 $\pm$ 2.2	0	$e^{\pm}j\tau^{\pm}$	160	162.8 $\pm$ 3.5	0
4j $\mu^{\pm}$	1363	1350.1 $\pm$ 37.7	0	2be $^{\pm}j\bar{p}$	19	19.4 $\pm$ 2.2	0	$e^{\pm}j\bar{p}\tau^{\mp}$	48	44.6 $\pm$ 3.3	0
3j high- Σp_T	159926	159143 $\pm$ 1061.9	0	2be $^{\pm}j$	63	63 $\pm$ 3.4	0	$e^{\pm}j\bar{p}\tau^{\pm}$	11	8.3 $\pm$ 1.5	0
3j low- Σp_T	62681	64213.1 $\pm$ 496	0	2be $^{\pm}$	96	92.1 $\pm$ 4.1	0	$e^{\pm}j\bar{p}$	121431	121023 $\pm$ 747.6	0
3j2 γ	151	177.5 $\pm$ 7.1	0	$\tau^{\pm}\tau^{\mp}$	856	872.5 $\pm$ 19	0	$e^{\pm}j\gamma\bar{p}$	159	192.6 $\pm$ 10.9	0
3j $\tau^{\pm}$ high- Σp_T	68	76.9 $\pm$ 3	0	$\gamma\bar{p}$	3793	3770.7 $\pm$ 127.3	0	$e^{\pm}j\gamma$	1389	1368.9 $\pm$ 38.9	0
3j $\bar{p}$ high- Σp_T	1706	1899.4 $\pm$ 77.6	0	$\mu^{\pm}\tau^{\mp}$	381	440.9 $\pm$ 7.3	0	$e^{\pm}j\mu^{\mp}\bar{p}$	42	33 $\pm$ 2.9	0
3j $\bar{p}$ low- Σp_T	42	36.2 $\pm$ 5.7	0	$\mu^{\pm}\bar{p}\tau^{\mp}$	60	75.7 $\pm$ 3.4	0	$e^{\pm}j\mu^{\pm}\bar{p}$	16	9.2 $\pm$ 1.9	0
3j $\gamma\tau^{\pm}$	39	37.8 $\pm$ 3.6	0	$\mu^{\pm}\bar{p}\tau^{\pm}$	15	12 $\pm$ 2	0	$e^{\pm}j\mu^{\mp}$	62	63.8 $\pm$ 3.2	0
3j $\gamma\bar{p}$	204	249.8 $\pm$ 24.4	0	$\mu^{\pm}\bar{p}$	734290	734296 $\pm$ 4897.8	0	$e^{\pm}j\mu^{\pm}$	13	8.2 $\pm$ 2	0
3j γ	24639	24899.4 $\pm$ 372.4	0	$\mu^{\pm}\gamma$	475	469.8 $\pm$ 12.5	0	$e^{\pm}e^{\mp}4j$	148	159.1 $\pm$ 7	0
3j $\mu^{\pm}\bar{p}$	2884	2971.5 $\pm$ 52.1	0	$\mu^{\pm}\mu^{\mp}\bar{p}$	169	198.5 $\pm$ 8.2	0	$e^{\pm}e^{\mp}3j$	717	743.6 $\pm$ 24.4	0
3j $\mu^{\pm}\gamma\bar{p}$	10	3.6 $\pm$ 1.9	0	$\mu^{\pm}\mu^{\mp}\gamma$	83	60 $\pm$ 3.1	0	$e^{\pm}e^{\mp}2j\bar{p}$	32	41.4 $\pm$ 5.6	0
3j $\mu^{\pm}\gamma$	15	7.9 $\pm$ 2.9	0	$\mu^{\pm}\mu^{\mp}$	25283	25178.5 $\pm$ 86.5	0	$e^{\pm}e^{\mp}2j\gamma$	10	11.4 $\pm$ 2.9	0
3j $\mu^{\pm}\mu^{\mp}$	175	177.8 $\pm$ 16.2	0	j2 $\gamma\bar{p}$	36	30.4 $\pm$ 4.2	0	$e^{\pm}e^{\mp}2j$	3638	3566.8 $\pm$ 72	0
3j $\mu^{\pm}$	5032	4989.5 $\pm$ 108.9	0	j2 γ	1822	1813.2 $\pm$ 27.4	0	$e^{\pm}e^{\mp}\tau^{\pm}$	18	16.1 $\pm$ 1.7	0
3b2j	23	28.9 $\pm$ 4.7	0	j $\tau^{\pm}$ high- Σp_T	52	56.2 $\pm$ 2.5	0	$e^{\pm}e^{\mp}\bar{p}$	822	831.8 $\pm$ 13.6	0
3bj	82	82.6 $\pm$ 5.7	0	j $\tau^{\pm}\tau^{\mp}$	203	252.2 $\pm$ 8.7	0	$e^{\pm}e^{\mp}\gamma$	191	221.9 $\pm$ 5.1	0
3b	67	85.6 $\pm$ 7.7	0	j $\bar{p}$ high- Σp_T	4432	4431.7 $\pm$ 45.2	0	$e^{\pm}e^{\mp}j\bar{p}$	155	170.8 $\pm$ 12.4	0
2 $\tau^{\pm}$	498	512.7 $\pm$ 14.2	0	j $\gamma\tau^{\pm}$	526	476 $\pm$ 9.3	0	$e^{\pm}e^{\mp}j\gamma$	48	45 $\pm$ 3.9	0
2 $\gamma\bar{p}$	128	107.2 $\pm$ 6.9	0	j $\gamma\bar{p}$	1882	1791.9 $\pm$ 72.3	0	$e^{\pm}e^{\mp}j$	17903	18258.2 $\pm$ 204.4	0
2 γ	5548	5562.8 $\pm$ 40.5	0	j γ	103319	102124 $\pm$ 570.6	0	$e^{\pm}e^{\mp}$	98901	99086.9 $\pm$ 147.8	0
2j high- Σp_T	190773	190842 $\pm$ 781.2	0	j $\mu^{\pm}\tau^{\mp}$	71	98 $\pm$ 3.9	0	b6j	51	42.3 $\pm$ 3.8	0
2j low- Σp_T	165984	162530 $\pm$ 1581	0	j $\mu^{\pm}\tau^{\pm}$	15	12 $\pm$ 2	0	b5j	237	192.5 $\pm$ 7.1	0
2j2 $\tau^{\pm}$	22	40.6 $\pm$ 3.2	0	j $\mu^{\pm}\bar{p}\tau^{\mp}$	26	30.8 $\pm$ 2.6	0	b4j high- Σp_T	26	23.4 $\pm$ 2.6	0
2j2 $\gamma\bar{p}$	11	8 $\pm$ 2.4	0	j $\mu^{\pm}\bar{p}$	109081	108323 $\pm$ 707.7	0	b4j low- Σp_T	836	821.7 $\pm$ 15.9	0
2j2 γ	580	581 $\pm$ 13.7	0	j $\mu^{\pm}\gamma\bar{p}$	171	171.1 $\pm$ 31	0	b3j high- Σp_T	12081	12071 $\pm$ 84.1	0
2j $\tau^{\pm}$ high- Σp_T	96	114.6 $\pm$ 3.3	0	j $\mu^{\pm}\gamma$	152	190 $\pm$ 39.3	0	b3j low- Σp_T	2974	2873 $\pm$ 31	0

10 most discrepant variables


CDF Run II Preliminary (2.0 fb^{-1})

Final State	Data	Background	σ	σ_t
$be^\pm p$	690	817.7 ± 9.2	-4.3	-2.7
$\gamma\tau^\pm$	1371	1217.6 ± 13.3	$+4.0$	$+2.2$
$\mu^\pm\tau^\pm$	63	35.2 ± 2.8	$+3.7$	$+1.7$
$b2j p$ high- Σp_T	255	327.2 ± 8.9	-3.7	-1.7
$2j\tau^\pm$ low- Σp_T	574	670.3 ± 8.6	-3.6	-1.5
$3j\tau^\pm$ low- Σp_T	148	199.8 ± 5.2	-3.5	-1.4
$e^\pm p\tau^\pm$	36	17.2 ± 1.7	$+3.5$	$+1.4$
$2j\tau^\pm\tau^\mp$	33	62.1 ± 4.3	-3.5	-1.3
$e^\pm j$	741710	764832 ± 6447.2	-3.5	-1.3
$j2\tau^\pm$	105	150.8 ± 6.3	-3.4	-1.2




- General agreement is good
- Largest deviation even is a deficit
- No sign of new physics

“Typical” variable: $W p_T$



Largest excess: $\gamma\tau$



- Up to now no indications from this approach
- However should be followed not to miss anything
- Makes only sense to plan for LHC once data are there and detector and SM physics is well understood

Conclusions

- The LHC can search for many new physics channels
- As a general rule new particles can be found up to $2 - 3 \text{ TeV}$
- However many models are not well defined, so limits should not be taken literally in many cases