

Physics at the LHC

Lecture 6: Higgs Physics at the LHC

Klaus Mönig, Sven Moch

(klaus.moenig@desy.de, sven-olaf.moch@desy.de)



Wintersemester 2010/2011

Reminder: Higgs mechanism in the Standard Model

- One complex Higgs doublet $\Phi = \begin{pmatrix} \Phi_1 + i\Phi_2 \\ H + i\Phi_3 \end{pmatrix}$ with vacuum expectation value $\begin{pmatrix} 0 \\ v \end{pmatrix}$, $v = 246 \text{ GeV}$.
- Higgs potential $V(\Phi) = \lambda(\Phi^*\Phi - v^2/2)^2$
- Gauge couplings of Higgs doublet give gauge boson masses:

$$m_W^2 = \frac{g^2 v^2}{4} \quad m_Z^2 = \frac{g^2 v^2}{4 \cos^2 \theta_W}$$


- Φ_i get absorbed in longitudinal d.o.f. of W,Z
- One Higgs particle H remains from fluctuations around v
 \Rightarrow Higgs couplings to gauge bosons fixed by gauge boson mass
- Higgs mass $m_H^2 = 2\lambda v^2$ free parameter

- Higgs couplings to fermions: $c_f \left[\tilde{F}_l(\Phi + v) f_r + \tilde{f}_r(\Phi + v) F_l \right]$
 \Rightarrow Higgs coupling \propto mass
- Partial widths:

$$\begin{aligned}\Gamma(H \rightarrow f\bar{f}) &= \frac{N_c^{(f)} G_\mu}{4\sqrt{2}\pi} m_f^2(m_H) m_H (1 + \delta_{QCD}^{(f)}) \\ \Gamma(H \rightarrow VV) &= \frac{3G_\mu^2 m_Z^4}{16\pi^3} m_H R_V (m_V^2/m_H^2) \\ &\rightarrow 2(1) \frac{\sqrt{2}G_\mu}{32\pi} m_H^3 \quad [V = W(Z)]\end{aligned}$$

- Higgs couplings to photons and gluons via loops

Partial and total widths of a Standard Model Higgs



- Light Higgs: $b\bar{b}$ is dominant
- Heavy Higgs: Mostly WW and ZZ
- Higgs width large above WW threshold

How the Higgs creates mass





and how it creates itself






Limits on m_H


Theory:

- No useful lower limit
- Upper limit from WW scattering:
 - Polarisation vector for longitudinally polarised particles $\propto p^2$
 - WW scattering without Higgs violates unitarity at 1.2 TeV





- Cross section gets regularised by the Higgs $\Rightarrow m_H \lesssim 1 \text{ TeV}$

- Perturbativity and vacuum stability give Higgs-mass limit as a function of the cutoff parameter
- If the SM is the final theory up to the Planck scale $\Lambda \sim 10^{19} \text{ GeV} \Rightarrow m_H \sim 120 - 180 \text{ GeV}$





Experimental:

- Direct searches at LEP exclude a Higgs below 114 GeV
- This limit is valid for couplings significantly smaller than in the SM
- Analyses of electroweak precision data suggest $m_H < 200$ GeV
- These analyses are only valid in the Standard Model without additional particles coupling to the gauge bosons




Higgs production at hadron colliders




- gg fusion produces a Higgs and nothing else in the detector
- All other graphs have associated particles that help in the tagging

Higgs production cross section

LHC



Tevatron



- gg fusion dominates in both cases
- At LHC WW-fusion significant, rest small
- At Tevatron WW-fusion and W, Z bremsstrahlung relevant
- Total cross section large at LHC, factor 100 smaller at Tevatron

How to discover a signal?

Assume background is known from somewhere

Number of events follows a Poisson distribution

⇒ variance σ for mean value n : $\sigma = \sqrt{n}$

For a given mass expect $n_t = n_s + n_b$ events

Require that signal is $> 5\sigma$ above background (corresponds to a probability of 10^{-7} for a background fluctuation)

⇒ need significance $S = n_s / \sqrt{n_b} > 5$

Influence of detector resolution: Gaussian signal with variance σ_D over linear background: $S \propto 1 / \sqrt{\sigma_D}$

Dependence on Luminosity: $S \propto \sqrt{\mathcal{L}}$

Higgs searches at the Tevatron

Light Higgs:

- Main decay to $b\bar{b}$
- Main channel $gg \rightarrow H \rightarrow b\bar{b}$ hopeless
- Possible channels $WH \rightarrow \ell\nu b\bar{b}$, $ZH \rightarrow \ell\ell b\bar{b}$, $ZH \rightarrow \nu\nu b\bar{b}$,

Medium Higgs

- $gg \rightarrow WW \rightarrow \ell\nu\ell\nu$ becomes accessible
- In addition some signal from $WH \rightarrow \ell\nu\ell\nu + \dots$


Heavy Higgs ($m_H > 200$ GeV)

- No chance because cross section too low


Search for $gg \rightarrow WW \rightarrow \ell\nu\ell\nu$

- Many variables with low separation power
- E.g. leptons correlated because of Higgs spin ($=0$)
- Combined with multivariate techniques, here NN
- Small signal under huge bg, WW and Drell-Yan dominant

ΔR for lepton pairs





Neural Net output



Results

- Low mass region dominated by $WH \rightarrow \ell\nu b\bar{b}$ and $WH, ZH \rightarrow \text{MET } b\bar{b}$
- Higher masses only $H \rightarrow \ell\nu\ell\nu$
- Exclusion at low masses still around $2 - 3\sigma(SM)$
- At $158 \text{ GeV} < m_H < 175 \text{ GeV}$ SM-Higgs excluded!

CDF limit for the different channels



Higgs search at the LHC


Easiest channel: $H \rightarrow ZZ \rightarrow \ell^+ \ell^- \ell^+ \ell^-$

$\sigma \times BR \sim 10 \text{ fb}$ for $m_H = 130 \text{ GeV}$

Largest background $t\bar{t} \rightarrow WbW\bar{b} \rightarrow \ell\nu \ell\nu c\bar{\nu} c\bar{\nu}$: $\sigma \times BR = 1300 \text{ fb}$

Best handle: lepton isolation and b-tagging


Dominant background after selection: ZZ



Channel can be used for discovery for $130 \text{ GeV} < m_H < 600 \text{ GeV}$

Discovery channel for light Higgs: $H \rightarrow \gamma\gamma$

Main backgrounds:





Start with $S/B = 10^{-6}$

→ need to reduce background by at least a factor 1000

Strategy:


- Effective selection of isolated photons (calorimeter granularity)
- Very good mass resolution (calorimeter energy resolution)

Complication: Experiments have \sim 1 radiation length in front of tracker
 ↳ about 50% of photons convert before they reach calorimeter
 ↳ have to include converted photons to keep high efficiency




The final signal is small, however visible with few fb^{-1} for a light Higgs

These two channels are sufficient to discover the Higgs over the full mass range



The Higgs in the VV-fusion channel




W-propagator:

$$\frac{1}{t-m_W^2} \text{ with } t = (p - p')^2 \approx 4pp' \sin^2 \frac{\theta}{2}$$



- propagator \sim constant as long as $2p' \sin \frac{\theta}{2} \lesssim m_W$
- forward jets should be visible in the detector

Higgs is colour singlet

- colour connection between forward jet and proton remnant
- rapidity gaps between forward jets and Higgs-jets




Largest background is $t\bar{t}$



Can be separated with rapidity cuts


With transverse mass $M_T = \sqrt{(E_{T,\ell\ell} + E_{T,miss})^2 - (\vec{p}_{T,\ell\ell} - \vec{p}_{T,miss})^2}$
 a clean signal can be separated




$H \rightarrow \tau\tau$:

- τ s have large energy
- Can assume that neutrinos go in direction of visible decay products
- τ energy can be reconstructed from kinematic constraints
- This works only if the τ s are not back-to-back


This allows $H \rightarrow \tau\tau$ reconstruction in the presence of tagging jets



With fusion channel the Higgs discovery will be assured in significantly more channels



Minimum luminosity needed for a Higgs discovery/exclusion



Higgs properties

LHC has discovered a particle compatible with a Higgs, what can be measured?


Mass:

Modes with complete Higgs reconstruction ($H \rightarrow \gamma\gamma$, $H \rightarrow ZZ \rightarrow 4\ell$) allow mass measurement with 0.1% precision.


Spin:

Coupling Hvv forbidden if H has spin 1 and v is massless vector particle (e.g. g or γ) (angular momentum conservation and Pauli principle)


→ visibility of $H \rightarrow \gamma\gamma$ or $gg \rightarrow H$ excludes spin 1



If $H \rightarrow ZZ \rightarrow 4\ell$ is visible spin/CP can be obtained from decay angle distributions:





- Take $H \rightarrow ZZ \rightarrow 2e2\mu$
- Add CP odd coupling to SM coupling with strength $\tan \xi / m_V^2$
- Most backgrounds can be suppressed by cuts
- Can distinguish the extreme cases



The width of the Higgs


- For a light Higgs the width is much smaller than the detector resolution
- If $m_H > 2m_W$ the width gets much larger and can be measured





For $m_H > 200$ GeV a precision $< 10\%$ is possible

Higgs couplings

- $\sigma \times BR \propto \frac{\Gamma_{prod} \Gamma_{dec}}{\Gamma_H}$
- Ratios of production rates measure ratios of partial widths
- Can obtain ratios of decay widths with $> 10\%$ accuracy





- For absolute partial widths need additional assumptions.
- Precisions of couplings depend on assumptions
- Minimal assumption $\Gamma_V < \Gamma_V^{SM}$ $V = W, Z$
- Again precision $> 10 - 20\%$
- Better precision with additional assumptions





Higgses at 7 (8) TeV

- The Higgs cross section at 7 TeV is a factor 4 smaller than at 14 TeV (at 8 TeV is 30% larger than at 7 TeV)
- The expected luminosity is 1 – few fb^{-1} in 2011
- The Tevatron might run for 3 more years to discover the Higgs
- It thus important to check if LHC can compete at 7(8) TeV
- There is an option to run in 2012 and do the upgrade to 14 TeV in 2013



- 1 fb^{-1} is sufficient to exclude the Higgs between 130 and around 400 GeV
- With 5 fb^{-1} the Higgs can be discovered with 3σ from the LEP-bound to 550 GeV
- Combining the two experiments this might even be possible in 2011



The Higgs in supersymmetric models

SUSY needs at least two Higgs-doublets (H_1, H_2) to generate masses of down- and up-type particles

Physical particles:

$$\begin{aligned} h &= H_2 \cos \alpha - H_1 \sin \alpha \\ H &= H_2 \sin \alpha + H_1 \cos \alpha \\ A &\quad \text{CP - odd} \\ H^\pm &\quad \text{charged Higgses} \end{aligned}$$

Define $\tan \beta = \frac{v_2}{v_1} = \text{ratio of expectation values}$ ($v_1^2 + v_2^2 = v_{SM}^2$)

Born Formulae:

$$m_{h,H}^2 = \frac{1}{2} \left[m_A^2 + m_Z^2 \mp \sqrt{(m_A^2 + m_Z^2)^2 - 4m_A^2 m_Z^2 \cos^2 2\beta} \right]$$

$$m_h < m_Z$$

$$m_H > m_Z$$

$$m_{H^\pm}^2 = m_A^2 + m_W^2$$



$$\tan 2\alpha = \tan 2\beta \frac{m_A^2 + m_Z^2}{m_A^2 - m_Z^2} \quad \left(-\frac{\pi}{2} < \alpha < 0 < \beta < \frac{\pi}{2} \right)$$

Higgs sector described by two free parameters

However large radiative corrections:

- shift of m_h up to ~ 130 GeV
- prediction gets dependent on other SUSY parameters, especially on mixing in stop sector
- strong dependence on top mass: $\Delta m_h / \Delta m_t \approx 1$

Currently allowed region:



$\tan\beta > 2$ preferred!

If m_A large:

- $\beta - \alpha = \pi/2$
 - $m_H \approx m_{H^\pm} \approx m_A$
- ⇒ Only one light Higgs can be seen

Branching ratios:

$$\Gamma(h \rightarrow VV) = \sin^2(\beta - \alpha)\Gamma_{\text{SM}}(h \rightarrow VV)$$
$$\Gamma(h \rightarrow U\overline{U}) = \frac{\cos^2 \alpha}{\sin^2 \beta}\Gamma_{\text{SM}}(h \rightarrow U\overline{U})$$
$$\Gamma(h \rightarrow D\overline{D}) = \frac{\sin^2 \alpha}{\cos^2 \beta}\Gamma_{\text{SM}}(h \rightarrow D\overline{D})$$

- For m_A large the light Higgs becomes SM like

Heavy Higgses at the LHC

For small $\tan \beta$ (disfavoured by LEP)

- $gg \rightarrow H, A$ production gets enhanced due to stronger $t\bar{t}H$ coupling
- $H, A \rightarrow t\bar{t}$ gets enhanced


For large $\tan \beta$

- H, A production from $b\bar{b}$ -fusion gets enhanced
- Large branching ratio $H \rightarrow \tau\tau$


Medium $\tan \beta$

- Only SM-like h visible

Visible SUSY-Higgs channels



Visible SUSY-Higgses at the LHC




- At least a SM-like h can always be seen
- However there is a significant chance that the rest of the SUSY Higgs sector remains invisible
- This does not include decays into SUSY particles (strongly model dependent!)


Invisible Higgs

- E.g. in SUSY models it is possible that the Higgs decays into invisible particles (e.g. the LSP)

- The properties of the VV-fusion process allows even in this case to separate the Higgs from the background



This allows to set a limit on $\xi^2 = BR(H \rightarrow \text{inv})\sigma(qq \rightarrow qqH)/\sigma(qq \rightarrow qqH, SM)$) in the 0.2-0.5 range



The Higgs thus cannot be missed because it decays invisible!

Conclusions of lecture 6

- A roughly SM like Higgs cannot be missed at the LHC
- A precise mass measurement is almost included in the discovery
- Spin and CP properties may be measured to some extend
- In SUSY the light h will be seen, the rest is questionable
- With a bit of luck we'll find the Higgs next year