

# Quantum Chromodynamics

## *lecture I*

**Sven-Olaf Moch**

*Universität Hamburg & DESY, Zeuthen*

---

*Belgian Dutch German summer school (BND 2012), Bonn, Sep 21, 2012*

# Plan

- *Introduction to QCD*

*Friday, September 21, 2012*

- QCD at work: infrared safety, factorization and evolution

*Saturday, September 22, 2012*

- Higgs boson production

*Sunday, September 23, 2012*

- Gauge boson production and QCD jets

*Monday, September 24, 2012*

- Top quark production

*Tuesday, September 25, 2012*

# Quantum Chromodynamics


- Fundamental forces: quantum fields with gauge symmetries
- Strong interaction: color- $SU(3)$ 
  - quarks (antiquarks): 6 spin-1/2 flavors in 3 colors

$$\begin{bmatrix} u_B \\ u_G \\ u_R \end{bmatrix} \quad \begin{bmatrix} d_B \\ d_G \\ d_R \end{bmatrix}, \quad \begin{bmatrix} s_B \\ s_G \\ s_R \end{bmatrix} \quad \begin{bmatrix} c_B \\ c_G \\ c_R \end{bmatrix}, \quad \begin{bmatrix} b_B \\ b_G \\ b_R \end{bmatrix} \quad \begin{bmatrix} t_B \\ t_G \\ t_R \end{bmatrix}$$

- gluons: 8 spin-1 color-anticolor-combinations

$$g_{B\bar{G}}, g_{R\bar{B}}, g_{G\bar{R}}, \dots, g_{B\bar{B}-G\bar{G}}, g_{B\bar{B}+G\bar{G}-2R\bar{R}}$$

- Interactions: Feynman diagrams



# QCD Lagrangian

- Classical part of QCD Lagrangian

$$\mathcal{L}_{\text{cl}} = -\frac{1}{4} F_{\mu\nu}^a F_a^{\mu\nu} + \sum_{\text{flavors}} \bar{\psi}_i (i\not{D} - m_q)_{ij} \psi_j$$

- Matter fields  $\psi_i, \bar{\psi}_j$  with  $i, j = 1, \dots, 3$  (fundamental rep.)
  - covariant derivative  $D_{\mu,ij} = \partial_\mu \delta_{ij} + ig_s (t_a)_{ij} A_\mu^a$
- Field strength tensor  $F_{\mu\nu}^a$  with  $a = 1, \dots, 8$  (adjoint rep.)
  - covariant derivative  $D_{\mu,ab} = \partial_\mu \delta_{ab} - g_s f_{abc} A_\mu^c$
  - $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g_s f_{abc} A_\mu^b A_\nu^c$
- Formal parameters of the theory (no observables)
  - strong coupling  $\alpha_s = g_s^2 / (4\pi)$
  - quark masses  $m_q$

## Quantization

- Gauge fixing (Feynman gauge  $\lambda = 1$ )  $\mathcal{L}_{\text{gauge-fix}} = -\frac{1}{2\lambda} (\partial^\mu A_\mu^a)^2$
- Ghosts (Grassmann fields  $\eta$ )  $\mathcal{L}_{\text{ghost}} = \partial_\mu \eta^{a\dagger} (D_{ab}^\mu \eta^b)$   
(removal of unphysical degrees of freedom for gauge fields) Fadeev, Popov

# From Lagrangian to Feynman rules

- Consider action  $S$

$$S = i \int d^4x (\mathcal{L}_{\text{cl}} + \mathcal{L}_{\text{gauge-fix}} + \mathcal{L}_{\text{ghost}}) = S_{\text{free}} + S_{\text{int}}$$

- Decompose action into free  $S_{\text{free}}$  and interacting part  $S_{\text{int}}$ 
  - $S_{\text{free}}$  contains bi-linear terms in fields
  - $S_{\text{int}}$  contains interactions
- Derivation of Feynman rules
  - inverse propagators from  $S_{\text{free}}$
  - interacting parts from  $S_{\text{int}}$  (in perturbative expansion)

# From Lagrangian to Feynman rules

- Consider action  $S$

$$S = i \int d^4x (\mathcal{L}_{\text{cl}} + \mathcal{L}_{\text{gauge-fix}} + \mathcal{L}_{\text{ghost}}) = S_{\text{free}} + S_{\text{int}}$$

- Decompose action into free  $S_{\text{free}}$  and interacting part  $S_{\text{int}}$ 
  - $S_{\text{free}}$  contains bi-linear terms in fields
  - $S_{\text{int}}$  contains interactions
- Derivation of Feynman rules
  - inverse propagators from  $S_{\text{free}}$
  - interacting parts from  $S_{\text{int}}$  (in perturbative expansion)

## Examples (I)

- Fermion propagator in QCD from  $\bar{\psi}_i \delta_{ij} (i\not{\partial} - m_q) \psi_j$ 
  - substitution  $\partial_\mu = -ip_\mu$  (Fourier transformation)
- Inverse propagator (momentum space)  $\Gamma_{ij}^{\bar{\psi}\psi}(p) = -i \delta_{ij} (\not{p} - m_q)$
- Check: quark propagator  $\Delta_{ij}(p) = +i \delta_{ij} \frac{1}{\not{p} - m_q + i0}$ 
  - causality in Minkowski space: prescription  $+i0$

## Examples (II)

- Gluon propagator in QCD from bi-linear terms in  $F_{\mu\nu}^a F_a^{\mu\nu}$  and  $\mathcal{L}_{\text{gauge-fix}}$

- recall  $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g_s f_{abc} A_\mu^b A_\nu^c$

- recall  $\mathcal{L}_{\text{gauge-fix}} = -\frac{1}{2\lambda} (\partial^\mu A_\mu^a)^2$

- Inverse propagator (momentum space)

$$\Gamma_{ab;\mu\nu}^{AA}(p) = +i \delta_{ab} \left[ p^2 g_{\mu\nu} - \left(1 - \frac{1}{\lambda}\right) p_\mu p_\nu \right]$$

- Gluon propagator  $\Delta^{ab;\mu\nu}(p) = +i \delta_{ab} \left[ \frac{-g_{\mu\nu}}{p^2} + (1 - \lambda) \frac{p_\mu p_\nu}{p^2} \right]$

- Check:  $\Gamma_{ac;\mu\rho}^{AA}(p) \Delta^{cb;\rho\nu}(p) = \delta_a^b g_\mu^\nu$

## Examples (II)

- Gluon propagator in QCD from bi-linear terms in  $F_{\mu\nu}^a F_a^{\mu\nu}$  and  $\mathcal{L}_{\text{gauge-fix}}$ 
  - recall  $F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a - g_s f_{abc} A_\mu^b A_\nu^c$
  - recall  $\mathcal{L}_{\text{gauge-fix}} = -\frac{1}{2\lambda} (\partial^\mu A_\mu^a)^2$
- Inverse propagator (momentum space)
$$\Gamma_{ab;\mu\nu}^{AA}(p) = +i \delta_{ab} \left[ p^2 g_{\mu\nu} - \left(1 - \frac{1}{\lambda}\right) p_\mu p_\nu \right]$$
- Gluon propagator  $\Delta^{ab;\mu\nu}(p) = +i \delta_{ab} \left[ \frac{-g_{\mu\nu}}{p^2} + (1 - \lambda) \frac{p_\mu p_\nu}{p^2} \right]$ 
  - Check:  $\Gamma_{ac;\mu\rho}^{AA}(p) \Delta^{cb;\rho\nu}(p) = \delta_a^b g_\mu^\nu$


## Examples (III)

- Interactions derived from  $S_{\text{int}}$ 
  - fermion-gluon interaction from  $\bar{\psi}_i i A_{ij} \psi_j \longrightarrow -i t_{ij}^a \gamma_\mu$
- General rule
  - replacement of all  $\partial_\mu$  by momenta  $p_\mu$   
(tedious for 3- and 4-gluon interactions)




# Feynman rules (I)


- Propagators
  - fermions, gluons, ghosts
  - covariant gauge



$$\delta^{ij} \frac{i}{\not{p} - m}$$




$$\delta^{ab} i \left( \frac{-g^{\mu\nu}}{p^2} + (1 - \lambda) \frac{p^\mu p^\nu}{(p^2)^2} \right)$$




$$\delta^{ab} \frac{i}{p^2}$$

# Feynman rules (II)


- Vertices



$$-i g (t^a)_{ji} \gamma^\mu$$




$$-g f^{abc} ((p - q)^\rho g^{\mu\nu} + (q - r)^\mu g^{\nu\rho} + (r - p)^\nu g^{\mu\rho})$$



$$-i g^2 f^{xac} f^{xbd} (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\sigma} g^{\nu\rho})$$

$$-i g^2 f^{xad} f^{xbc} (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\rho} g^{\nu\sigma})$$

$$-i g^2 f^{xab} f^{xcd} (g^{\mu\rho} g^{\nu\sigma} - g^{\mu\sigma} g^{\nu\rho})$$



$$g f^{abc} q^\mu$$


# Color algebra

- $SU(N)$ -generators  $t^a$  from fundamental representation

$$\text{Tr} \left( t^a t^b \right) = \frac{1}{2} \delta^{ab}$$


- $SU(N)$ -generators  $f^{abc}$  of adjoint representation

$$f^{abc} = i \text{Tr} \left( \left[ t^a, t^b \right] t^c \right)$$



- Fierz identity

$$(t^a)_{i_1}^{j_1} (t^b)_{i_2}^{j_2} = \delta_{i_1}^{j_2} \delta_{i_2}^{j_1} - \frac{1}{N} \delta_{i_1}^{j_1} \delta_{i_2}^{j_2}$$



## Useful relations

- Color constant (quadratic Casimir)

- quarks  $(t^a t^a)_{ij} = C_F \delta_{ij} = \frac{N^2 - 1}{2N} \delta_{ij} = \frac{4}{3} \delta_{ij}$

- gluons  $f^{acd} f^{bcd} = C_A \delta_{ab} = N \delta_{ab} = 3 \delta_{ab}$

# Renormalization

## Physics picture

- Parameters of Lagrangian in quantum field theory have no unique physical interpretation
- Generic quantity  $R$  depends on
  - hard scale  $Q$ , mass  $m_q$
  - in perturbative study on coupling constant  $\alpha_s$
- Radiative corrections
  - resolve quantum fluctuations at given resolution length  $a \sim 1/\mu$
  - induce dependence of  $R$  on scale  $\mu$

# Renormalization

## Physics picture



- Parameters of Lagrangian in quantum field theory have no unique physical interpretation
- Generic quantity  $R$  depends on
  - hard scale  $Q$ , mass  $m_q$
  - in perturbative study on coupling constant  $\alpha_s$
- Radiative corrections
  - resolve quantum fluctuations at given resolution length  $a \sim 1/\mu$
  - induce dependence of  $R$  on scale  $\mu$
- Renormalization “group” governed by QCD describes changes  $R$  with respect to  $\mu$  (differential equation of first order)

$$\left\{ \mu^2 \frac{\partial}{\partial \mu^2} + \beta(\alpha_s) \frac{\partial}{\partial \alpha_s} - \gamma_m(\alpha_s) m_q \frac{\partial}{\partial m_q} \right\} R \left( \frac{Q^2}{\mu^2}, \alpha_s, \frac{m_q^2}{Q^2} \right) = 0$$



- partial derivatives  $\beta(\alpha_s) = \frac{\partial}{\partial \mu^2} \alpha_s$  and  $\gamma_m(\alpha_s) m_q = \frac{\partial}{\partial \mu^2} m_q$
- solution of differential equation requires initial conditions  
→ definition of renormalization scheme

# Running coupling

- Effective coupling constant  $\alpha_s$  depends on resolution
- QCD distinguished by self-interaction of gluons; e.g. vertex corrections





– screening (like in QED)





– anti-screening (color charge of  $g$ )

# Running coupling

- Effective coupling constant  $\alpha_s$  depends on resolution
- QCD distinguished by self-interaction of gluons; e.g. vertex corrections



– screening (like in QED)



– anti-screening (color charge of  $g$ )

- Scale dependence governed by  $\beta$ -function of QCD



$$\mu^2 \frac{d}{d\mu^2} \alpha_s(\mu) = \beta(\alpha_s) = -\beta_0 \alpha_s^2 - \beta_1 \alpha_s^3 - \beta_2 \alpha_s^4 - \beta_3 \alpha_s^5 - \dots$$

- QCD  $\beta$ -function has negative sign
- perturbative expansion with coefficients  $\beta_0, \beta_1, \beta_2, \dots$

$$\beta_0 = \frac{1}{4\pi} \left( \frac{11}{3} C_A - \frac{2}{3} n_f \right) = \frac{1}{4\pi} (7) \quad (\text{for } n_f = 6)$$

# Asymptotic freedom

- Solution of QCD  $\beta$ -function
  - perturbative expansion to four loops *van Ritbergen, Vermaseren, Larin '97*
  - very good convergence of perturbative series even at low scales (but  $\alpha_s \gg \alpha_{\text{QED}}$ )






# Strong coupling constant

## Essential facts

- $\alpha_s(M_Z)$  from  $e^+e^-$  data high
- $\alpha_s(M_Z)$  from DIS data low
- World average 1992  
 $\alpha_s(M_Z) = 0.117 \pm 0.004$

Process	Ref.	Q [GeV]	$\alpha_s(Q)$	$\alpha_s(M_{Z^0})$	$\Delta\alpha_s(M_{Z^0})$		order of perturb.
					exp.	theor.	
1 $R_\tau$ [LEP]	[7-10]	1.78	$0.318 \pm^{0.048}_{0.039}$	$0.117 \pm^{0.006}_{0.005}$	$\pm^{0.003}_{0.004}$	$\pm^{0.005}_{-0.004}$	NNLO
2 $R_\tau$ [world]	[2]	1.78	$0.32 \pm 0.04$	$0.118 \pm^{0.004}_{0.006}$	-	-	NNLO
3 DIS [ $\nu$ ]	[3]	5.0	$0.193 \pm^{0.019}_{0.018}$	$0.111 \pm^{0.006}_{0.007}$	$\pm^{0.004}_{-0.006}$	0.004	NLO
4 DIS [ $\mu$ ]	[12]	7.1	$0.180 \pm 0.014$	$0.113 \pm 0.005$	0.003	0.004	NLO
5 $J/\Psi, \Upsilon$ decay	[4]	10.0	$0.167 \pm^{0.015}_{0.011}$	$0.113 \pm^{0.007}_{0.005}$	-	-	NLO
6 $e^+e^-$ [ $\sigma_{had}$ ]	[14]	34.0	$0.163 \pm 0.022$	$0.135 \pm 0.015$	-	-	NNLO
7 $e^+e^-$ [shapes]	[15]	35.0	$0.14 \pm 0.02$	$0.119 \pm 0.014$	-	-	NLO
8 $p\bar{p} \rightarrow b\bar{b}X$	[11]	20.0	$0.136 \pm^{0.025}_{0.024}$	$0.108 \pm^{0.015}_{0.014}$	0.006	$\pm^{0.014}_{-0.013}$	NLO
9 $p\bar{p} \rightarrow W$ jets	[13]	80.6	$0.123 \pm 0.027$	$0.121 \pm 0.026$	0.018	0.020	NLO
10 $\Gamma(Z^0 \rightarrow had.)$	[5]	91.2	$0.133 \pm 0.012$	$0.133 \pm 0.012$	0.012	$\pm^{0.003}_{-0.001}$	NNLO
11 $Z^0$ ev. shapes							
ALEPH	[7]	91.2	$0.119 \pm^{0.008}_{0.010}$		-	-	NLO
DELPHI	[8]	91.2	$0.113 \pm 0.007$		0.002	0.007	NLO
L3	[9]	91.2	$0.118 \pm 0.010$		-	-	NLO
OPAL	[10]	91.2	$0.122 \pm^{0.006}_{0.005}$		0.001	$\pm^{0.006}_{-0.005}$	NLO
SLD	[6]	91.2	$0.120 \pm^{0.015}_{0.013}$		0.009	$\pm^{0.012}_{-0.009}$	NLO
Average	[6-10]	91.2		$0.119 \pm 0.006$	0.001	0.006	NLO
12 $Z^0$ ev. shapes							
ALEPH	[7]	91.2	$0.125 \pm 0.005$		0.002	0.004	resum.
DELPHI	[8]	91.2	$0.122 \pm 0.006$		0.002	0.006	resum.
L3	[9]	91.2	$0.126 \pm 0.009$		0.003	0.008	resum.
OPAL	[10]	91.2	$0.122 \pm^{0.003}_{-0.006}$		0.001	$\pm^{0.003}_{-0.006}$	resum.
Average	[7-10]	91.2		$0.123 \pm 0.005$	0.001	0.005	resum.

Table 1: Summary of measurements of  $\alpha_s$ . For details see text.

World Summary of  $\alpha_s$  2012:

$$\rightarrow \alpha_s(M_Z) = 0.1185 \pm 0.0007$$

(preliminary)

S. Bethke (Karlsruhe 2012)


$$\Lambda_{\overline{\text{MS}}}^{(5)} = (214 \pm 9) \text{ MeV}$$

$$\Lambda_{\overline{\text{MS}}}^{(4)} = (297 \pm 11) \text{ MeV}$$

# Renormalization

## Technicalities


- Radiative corrections require integration over loop momenta
  - loop integrals can diverge in ultraviolet  $l \rightarrow \infty$
  - power counting reveals divergence in ultraviolet
- Example: self-energy in scalar field theory (off-shell momentum  $q^2 \neq 0$ )

$$\int d^4l \frac{1}{l^2(l-q)^2}$$



# Renormalization

## Technicalities

- Radiative corrections require integration over loop momenta
  - loop integrals can diverge in ultraviolet  $l \rightarrow \infty$
  - power counting reveals divergence in ultraviolet
- Example: self-energy in scalar field theory (off-shell momentum  $q^2 \neq 0$ )

$$\int d^4l \frac{1}{l^2(l-q)^2}$$


Nobel prize 1999



*The formulas of 't Hooft and Veltman made the infinities vanish into thin air!*

*The contributions of the Nobel Laureates seen as a "mathematical machinery".*

### Goodbye to infinities

For decades, attempts were made to explain the weak interactions. But meaningless results often appeared in the form of infinite probabilities and infinite so-called quantum corrections.


't Hooft and Veltman showed how these nasty infinities could be tamed and interpreted. In their "mathematical machinery" the theory is first modified, among other things, through the introduction of a number of "ghost particles". Calculations are then run in an unreal space-time in which the number of dimensions is a shade lower than the real num-

Quantum Chromodynamics – p.15

# Regularization

- Example for UV divergent loop integral (Euclidean region):

$$\int \frac{d^4 l}{(2\pi)^4} \frac{1}{l^2 + m_q^2}$$



- Various regularization methods

- cut-off  $l \leq \Lambda_{\text{cut-off}}$ ; Pauli-Villars  $\frac{1}{l^2} \rightarrow \frac{1}{l^2 - M^2}$ ; lattice; ...

## Dimensional regularization

- Lorentz invariance and  $SU(N)$  gauge invariance manifest
- Analytical continuation in space-time dimension  $D = 4 - 2\epsilon$

- loop integral  $\int \frac{d^4 l}{(2\pi)^4} \rightarrow \int \frac{d^D l}{(2\pi)^D}$
- Lorentz index  $\mu \in \{0, 1, 2, 3\} \rightarrow \{0, 1, \dots, D\}$
- Lorentz vector  $p^\mu \in (p^0, p^1, p^2, p^3) \rightarrow (p^0, p^1, \dots, p^{D-1})$
- metric  $g^{\mu\nu} g_{\mu\nu} = g_\mu^\mu = D$
- Dirac algebra  $\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}$  and  $\gamma^\mu \gamma^\nu \gamma_\mu = (2 - D)\gamma^\nu$

# Renormalization

## In a nut-shell

- compute vertex corrections (one-particle irreducible diagrams)





- three-gluon vertex

$$V_{A_\mu A_\nu A_\rho} = g_s A_\mu A_\nu A_\rho$$

- gluon-quark vertex

$$V_{\psi\bar{\psi}A_\mu} = g_s A_\mu \bar{\psi}\psi$$

- compute self-energy corrections



- Redefinition of fields, vertices and parameters in Lagrangian

- vertices  $V_{A_\mu A_\nu A_\rho}^b = Z_1 V_{A_\mu A_\nu A_\rho}^r$  and  $V_{\psi\bar{\psi}A_\mu}^b = Z_{1F} V_{\psi\bar{\psi}A_\mu}^r$

- fields  $\psi^b = (Z_2)^{1/2} \psi^r$  and  $A_\mu^b = (Z_3)^{1/2} A_\mu^r$

- parameters coupling constant  $g_s^b = Z_g g_s^r$  and mass  $m^b = Z_m m^r$

# Renormalization

## Gauge invariance

- Renormalization of vertex corrections imply

$$Z_1 = Z_g (Z_3)^{3/2}, \quad Z_{1F} = Z_g (Z_3)^{1/2} Z_2$$

- Combinations of  $Z$ -factors fixed by  $SU(N)$  gauge invariance of QCD
  - Slavnov-Taylor (or Ward) identities

$$Z_g (Z_3)^{1/2} = \frac{Z_1}{Z_3} = \frac{Z_{1F}}{Z_2}$$

## Renormalized Lagrangian

- Construction of renormalized QCD Lagrangian  $\mathcal{L}^{\text{ren}}$  with rescaled fields, parameters. etc.
- $\mathcal{L}^{\text{bare}}$  decomposed into  $\mathcal{L}^{\text{ren}}$  and counter term  $\mathcal{L}^{\text{ct}}$

$$\mathcal{L}^{\text{bare}}(\psi^{\text{b}}, \bar{\psi}^{\text{b}}, A_\mu^{\text{b}}) = \mathcal{L}^{\text{ren}}(\psi^{\text{r}}, \bar{\psi}^{\text{r}}, A_\mu^{\text{r}}) + \mathcal{L}^{\text{ct}}$$

- $\mathcal{L}^{\text{ct}}$  contains all parameters with factors  $(Z_i - 1)$
- ultraviolet divergences absorbed by  $\mathcal{L}^{\text{ct}}$


# Quark mass renormalization

- Heavy quark self-energy  $\Sigma(p, m_q)$

$$\longrightarrow + \longrightarrow \textcircled{\Sigma} \longrightarrow + \longrightarrow \textcircled{\Sigma} \textcircled{\Sigma} \longrightarrow + \dots = \frac{i}{\not{p} - m_q - \Sigma(p, m_q)}$$

## QCD

- QCD corrections to self-energy  $\Sigma(p, m_q)$ 
  - dimensional regularization  $D = 4 - 2\epsilon$
  - one-loop: UV divergence  $1/\epsilon$  (Laurent expansion)



$$\Sigma^{(1), \text{bare}}(p, m_q) = \frac{\alpha_s}{4\pi} \left( \frac{\mu^2}{m_q^2} \right)^\epsilon \left\{ (\not{p} - m_q) \left( -C_F \frac{1}{\epsilon} + \text{fin.} \right) + m_q \left( 3C_F \frac{1}{\epsilon} + \text{fin.} \right) \right\}$$

- Relate bare and renormalized mass parameter  $m_q^{\text{bare}} = m_q^{\text{ren}} + \delta m_q$

$$\textcircled{\Sigma^{\text{ren}}(p, m_q)} = \longrightarrow + \longrightarrow \textcircled{\text{gluon loop}} \longrightarrow + \longrightarrow \times \longrightarrow + \dots$$

$$(Z_\psi - 1)\not{p} - (Z_m - 1)m_q$$



# Mass renormalization scheme

## Pole mass

- Based on (unphysical) concept of top quark being a free parton
  - $m_q^{\text{ren}}$  coincides with pole of propagator at each order

$$\not{p} - m_q - \Sigma(p, m_q) \Big|_{\not{p}=m_q} \rightarrow \not{p} - m_q^{\text{pole}}$$

- Definition of pole mass ambiguous up to corrections  $\mathcal{O}(\Lambda_{QCD})$ 
  - heavy quark self-energy  $\Sigma(p, m_q)$  receives contributions from regions of all loop momenta – also from momenta of  $\mathcal{O}(\Lambda_{QCD})$
  - bound from lattice QCD:  $\Delta m_q \geq 0.7 \cdot \Lambda_{QCD} \simeq 200 \text{ MeV}$   
Bauer, Bali, Pineda '11

## $\overline{MS}$ scheme

- $\overline{MS}$  mass definition
  - one-loop minimal subtraction

$$\delta m_q^{(1)} = m_q \frac{\alpha_s}{4\pi} 3C_F \left( \frac{1}{\epsilon} - \gamma_E + \ln 4\pi \right)$$

- $\overline{MS}$  scheme induces scale dependence:  $m(\mu)$

# Running quark mass


## Scale dependence

- Renormalization group equation for scale dependence
  - mass anomalous dimension  $\gamma$  known to four loops

Chetyrkin '97; Larin, van Ritbergen, Vermaseren '97

$$\left( \mu^2 \frac{\partial}{\partial \mu^2} + \beta(\alpha_s) \frac{\partial}{\partial \alpha_s} \right) m(\mu) = \gamma(\alpha_s) m(\mu)$$

- Plot mass ratio  $m_t(163\text{GeV})/m_t(\mu)$



# Scheme transformations

- Conversion between different renormalization schemes possible in perturbation theory
- Relation for pole mass and  $\overline{MS}$  mass
  - known to three loops in QCD Gray, Broadhurst, Gräfe, Schilcher '90; Chetyrkin, Steinhauser '99; Melnikov, v. Ritbergen '99
  - EW sector known to  $\mathcal{O}(\alpha_{EW}\alpha_s)$  Jegerlehner, Kalmykov '04; Eiras, Steinhauser '06
  - example: one-loop QCD

$$m^{\text{pole}} = m(\mu) \left\{ 1 + \frac{\alpha_s(\mu)}{4\pi} \left( \frac{4}{3} + \ln \left( \frac{\mu^2}{m(\mu)^2} \right) \right) + \dots \right\}$$

# Treatment of heavy quarks

- Light quarks:  $m_u, m_d \ll \Lambda_{\text{QCD}}, \quad m_s < \Lambda_{\text{QCD}}$ 
  - neglect “light quark” masses in hard scattering process
- Heavy quarks:  $m_c, m_b, m_t \gg \Lambda_{\text{QCD}}$ 
  - mass effects important

## Example

- Different kinematical regions for  $m_c$ 
  - $Q \not\gg m_c$ : partons  $u, d, s, g$  with  $n_f = 3$   
massive charm quark and terms  $m_c/Q \neq 0$  are sizable
  - $Q \gg m_c$ : partons  $u, d, s, c, g$  with  $n_f = 4$   
massless charm quark and terms  $m_c/Q \rightarrow 0$  are neglected

# Decoupling

## *In a nut-shell*

- QCD with different number of quarks can be related  $\longrightarrow$  matching of two distinct theories
- Heavy quarks can be decoupled in limit  $m_q \rightarrow \infty$  Appelquist, Carrazzone '74
- Consider QCD parameters in both theories and match at scale  $\mu$ 
  - $n_l$  light flavors +  $n_h$  heavy quarks of masses  $m_q$  at low scales
  - $n_l + n_h$  light flavors at high scales
- Example: running coupling constant

$$\alpha_s^{n_l} \longrightarrow \alpha_s^{(n_l+n_h)}$$

## $\overline{\text{MS}}$ scheme


- Decoupling theorem in the  $\overline{\text{MS}}$ -scheme not true in naive sense
  - mass effects not  $1/m_q$  suppressed in theory with  $n_l$  light and  $n_h$  heavy flavors
  - anomalous dimensions exhibit discontinuities at flavor thresholds
- Decoupling constants in the  $\overline{\text{MS}}$  scheme

Larin, van Ritbergen, Vermaseren '94; Chetyrkin, Kniehl, Steinhauser '97

# $\alpha_s$ with flavor thresholds


- Solution of QCD  $\beta$ -function
  - discontinuities for  $n_f = 3 \rightarrow n_f = 4 \rightarrow n_f = 5$
- Big picture

Bethke for PDG 2012



# $\alpha_s$ with flavor thresholds

- Solution of QCD  $\beta$ -function
  - discontinuities for  $n_f = 3 \rightarrow n_f = 4 \rightarrow n_f = 5$
- Zoom



# Summary (part I)

## *QCD: the gauge theory of the strong interaction*

- Quarks and gluons as classical degrees of freedom
- Quantum corrections determine dynamical properties
  - scale dependence of observables
  - running coupling constant and asymptotic freedom
- Renormalization required by quantum corrections
  - subtraction of ultraviolet singularities
  - definition of renormalization scheme
  - parameters of Lagrangian are not observables  $\alpha_s, m_q, \dots$